



# Birds of the Ramsar site Estação Ecológica de Taiamã and buffer zone, Pantanal wetlands, Brazil

Angélica Vilas Boas da Frota, Breno Dias Vitorino, Carolina Joana da Silva, Solange Kimie Ikeda-Castrillon, Josué Ribeiro da Silva Nunes

Programa de Pós-graduação em Ciências Ambientais, Universidade do Estado de Mato Grosso, Centro de Pesquisa em Limnologia, Etnobiologia e Biodiversidade do Pantanal, Av. Santos Dumont, s/n° - Cidade Universitária (Block II), Cáceres, 78200-000, MT, Brazil.

**Corresponding author:** Angélica Vilas Boas da Frota, [angelicafrotaa@gmail.com](mailto:angelicafrotaa@gmail.com)

## Abstract

The Estação Ecológica de Taiamã (EET) is one of the Ramsar sites in the Pantanal wetlands of Brazil. We present an annotated list of bird species recorded between 2015 and 2018, along with a compilation of existing data for this area. In our sampling, 207 species were recorded, with 76% documented. We have the first record of *Chlidonias niger* (Linnaeus, 1758) for the Pantanal and Central region of Brazil. Another 30 species are new records for EET, such as the migrant *Hirundo rustica* Linnaeus, 1758. The general list comprises 24 orders, 60 families, and 278 species, of which 160 were assigned to the primary list and 118 to the secondary list. This study will aid in the conservation of biodiversity in the Pantanal. The EET and its buffer zone are important for maintaining biodiversity and expanding its boundaries may further promote the conservation of birds in Brazilian wetlands.

## Keywords

Bird migration, floodplain, Nearctic migrant, ornithological inventory, Paraguay River, protected areas, waterbirds.

**Academic editor:** Rafael Antunes Dias | Received 7 June 2019 | Accepted 3 March 2020 | Published 10 April 2020

**Citation:** Frota AVB, Vitorino BD, da Silva CJ, Ikeda-Castrillon SK, Nunes JRS (2020) Birds of the Ramsar site Estação Ecológica de Taiamã and buffer zone, Pantanal wetlands, Brazil. Check List 16 (2): 401–422. <https://doi.org/10.15560/16.2.401>

## Introduction

The Pantanal wetland ecosystem is regulated by the natural flood pulse process, with its biodiversity adapted to water dynamics and the heterogeneous landscape (Junk et al. 1989; Junk and Da Silva 1996). In the Brazilian Pantanal, 582 bird species have been reported (Nunes 2011), of which 104 depend directly on wetlands (Junk et al. 2006). Bird studies in this floodplain still present gaps in the documented record, and a more systematic analysis of taxa, especially for the northwest region is required (Tubelis and Tomas 2003). Given the increasing human pressure on the Pantanal wetland ecosystem, such as large infrastructure projects, agricultural

expansion, mining, and the impacts of global climate change (Tomas et al. 2019), ornithological inventories provides important information for developing conservation strategies and actions for biodiversity.

In the Pantanal wetlands is the Ramsar site Estação Ecológica de Taiamã (EET), which has been a full protection area since 1981 by decree 86.061; the EET was assigned as a Wetland of International Importance in 2018 (Brasil 2019). The main objectives of the EET is to preserve nature and conduct scientific studies in accordance with Federal Law 9.985 of 2000 (Brasil 2000). Even with some known bird records (Lopes et al. 2016; Brasil 2017), there is difficulty with accessing

and conducting standardised studies in the EET due to its characteristics in the fluvio-lacustrine region of the upper Paraguay river (Wantzen et al. 2005).

Information contained in species lists is useful for understanding and conserving biodiversity and the environmental (Silveira et al. 2010). Biodiversity knowledge is essential for planning and taking action to protect Brazil's wetlands, and our research aimed to fill in the gaps in the database for this region's wetland birds. The objective of our study was to inventory and consolidate a list of bird species occurring on Ramsar site Estação Ecológica de Taiamã. We also discuss some aspects of the biology and conservation of these species.

## Methods


**Study area.** The EET is a protected river island in the upper Paraguay river basin, located at 16°48'S–16°58'S and 57°24'W–57°40'W in the Cáceres municipality, Mato Grosso, Brazil (Fig. 1). The EET includes 11,555 ha of Pantanal wetland between the Paraguay and Bracinho rivers. According to Kottek et al. (2006), the climate is Aw (hot and humid), with an annual rainfall of 1500 mm and annual average maximum and minimum temperatures of 32 °C and 20 °C, respectively. Veloso et al. (1991) classified the vegetation as Alluvial Semideciduous Seasonal Forest.

The EET is characterised by different functional units. These are Aquatic-Terrestrial Transition Zones (ATTZ), swamps, and permanently aquatic areas. Among

the main macrohabitats are the Purple Coraltree Monospecific Forest (forest dominated by *Erythrina fusca* Lour., which is locally known as “abobral”), Shrubland, and Pioneer Polyspecific Forest, which is present of the river margins, Flooded Grasslands composed of herbaceous plants, aquatic macrophytes, and floating islands and meadows (locally known as “campos de batumes”, composed of plants and layers of organic matter accumulated by hydrological changes), lakes, and river channels (Frota et al. 2017). The buffer zone is characterised by irregular depression lakes, some of which are interconnected by floodplain channels (Wantzen et al. 2005), flooded grasslands, aquatic macrophytes, abobral, and campos de batumes.

**Sampling and analysis.** We collected systematic and non-systematic data in various EET macrohabitats and the buffer zone. First, we sampled 36 field days of the 2015–2016 hydrological cycle. Subsequently, we carried out new expeditions in the area using the Long-Term Ecological Research (LTER) in the Pantanal wetland “Ecological Dynamics of the Upper Paraguay River floodplain” and added 40 field days from the 2017–2018 hydrological cycle to increase the composition of the species list. We identified bird species and counted individuals by visual and sound contact, using 10 × 42 mm binoculars, a digital recorder, photo cameras, bird guides (Erize et al. 2006; Ridgely and Tudor 2009; Gwynne et al. 2010) and specialised literature (Sick 1997; del Hoyo et al. 2017).

Whenever possible, we documented bird species



**Figure 1.** Estação Ecológica de Taiamã (EET), Pantanal wetland, Brazil: localization, delimitation and sampling points in the study area.

with photos and audio recordings, which were archived on Wikiaves (<http://www.wikiaves.com.br>). We followed the Brazilian Ornithological Records Committee's list (Piacentini et al. 2015) for taxonomic classification. We used the Brazilian Red Book of Threatened Species of Fauna (Brasil 2018) for the national conservation status and the Red List of Threatened Species (IUCN 2019) for the international conservation status. For migratory behaviour, we followed the classification of Somenzari et al. (2018): Migratory (MGT), Partially Migratory (MPR), Vagrant (VAG), and Not Defined (ND).

For the survey, we used point counts (Blondel et al. 1970) along the rivers with a 50 m fixed radius and 15 minutes to record all contacts. We sample three macrohabitats types (two stations for each) in the 2015–2016 cycle: Shrubland and Pioneer Forest, *E. fusca* Monospecific Forest and Flooded Grassland. We distributed sixteen points along each macrohabitat type, spaced at least 200 m from each other. Each point was sampled for three consecutive days in two seasons each (wet and dry), at dawn and dusk, totaling 72 hours. In the 2017–2018 cycle, we sampled five tracks in the LTER site and 10 points spaced at least 500 m from each other. The points were sampled at dawn for two days in four hydrological periods (flood, ebb, dry, and flooding), totalling 100 observation-hours. In total, 11 sites (Table 1) and 98 points were sampled on the smaller boat (see Appendix, Table S1).

We assigned the frequency of occurrence (FO) for each species as the number of samples where a given species was recorded divided by the total number of samples (D'Angelo Neto et al. 1998). The categories for FO were: very frequent (100–75%); frequent (74–50%); reasonably frequent (49–25%); less frequent (<25%), and infrequent for non-systematic data.

For the new record, we considered the species range maps of BirdLife International and Handbook of the Birds of the World (2018) on the cartographic base. We used QGIS v. 3.4.4 to draw a map with our new record. We also included secondary records for other places of South America (Dias et al. 2010; Quiroga et al. 2015; Gonsioroski; 2018; Franz et al. 2018; Wikiaves 2019),

and that are outside BirdLife's species range maps.

To compile the species list for the EET (systematic and non-systematic data), we combined our field data with previously existing records, including published information (Kantek and Onuma 2013; Lopes et al. 2016), specimens deposited in the scientific collection of Department of Zoology at the Universidade Federal de Minas Gerais (DZUFMG), and the bird list of Estação Ecológica de Taiamã's Management Plan (Brasil 2017).

For the general list of birds, each species was reviewed and separated into three lists according to the methodology adopted by Piacentini et al. (2015) and other check list studies (Nunes et al. 2017; Vitorino et al. 2018):

- Primary list: species with documented evidence for the area in the form of a complete or partial specimen, photograph, or sound recording.
- Secondary list: species with records of occurrence for the area without known documentary evidence. The species were considered of probable occurrence, based on the currently known distribution.
- Tertiary list: species with records of occurrence for the area without known documentary evidence; the occurrence is questionable, invalid taxonomy, or non-existent.

The general list is, thus, composed of species from the primary and secondary lists. The documentation type of ornithological records was based on Carlos et al. (2010), even as applied to another protected area in Brazil (Vitorino et al. 2018): "A-level documentation" for species with specimens deposited in scientific collections; "B-level documentation" for species documented with photographs or audio recordings; and "Indication of Occurrence" for species recorded in the field, but not documented.

## Results

We found 23 orders, 58 families, and 207 bird species occurring in the EET and buffer zone. Of these, 46 species were non-systematic and 31 were exclusive in the LTER expeditions. Of the total species recorded in the field, 76% have vouchers (Table 2). The most representative families

**Table 1.** List of sampling sites in the Estação Ecológica de Taiamã (EET), Pantanal wetland, Brazil, with sampled hydrological cycle, geographic coordinates (at center point of sites) and predominant macrohabitats.

Hydrological cycle	Site names	Site code	Latitude	Longitude	Macrohabitats
2015–2016	Station 1	S1	16°49.28'S	057°32.30'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
	Station 2	S2	16°51.95'S	057°32.26'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels
	Station 3	S3	16°51.92'S	057°32.88'W	Shrubland and pioneer forest / river channels
	Station 4	S4	16°52.48'S	057°26.29'W	Shrubland and pioneer forest / river channels
	Station 5	S5	16°50.04'S	057°38.14'W	Flooded grassland / river and floodplain channels
	Station 6	S6	16°52.51'S	057°24.45'W	Flooded grassland / lakes / river channels
2017–2018	Track 1	T1	16°49.53'S	057°31.13'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
	Track 2	T2	16°52.64'S	057°25.69'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
	Track 3	T3	16°54.06'S	057°28.63'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / river channels
	Track 4	T4	16°51.86'S	057°32.41'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / aquatic macrophytes / river channels
	Track 5	T5	16°50.06'S	057°39.70'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels

**Table 2.** List of birds recorded at Estação Ecológica de Taimã (EET), Pantanal wetland, Brazil. Legend: *Migratory status* (Somenzari et al. 2018): resident (RDT), migratory (MGT), partially migratory (MPR), vagrant (VAG), and not defined (ND). *Occurrence status of this study*: very frequent (100–75%), frequent (74–50%), reasonably frequent (49–25%), less frequent (<25%), and infrequent for non-systematic data (IN). *Record type* (Carlos et al. 2010): IO = indication of occurrence (species not documented), AD = A-level documentation (species with specimens deposited in scientific collections), and BD = B-level documentation (species documented with photographs or audio recordings). *Source*: this study (1), Lopes et al. 2016 (2), Kantek and Onuma 2013 (3), specimens deposited in the Department of Zoology of the Universidade Federal de Minas Gerais (4), and Brazil 2017 (5); Secondary list species in brackets. VU = Vulnerable (IUCN 2019) and VU\* = Vulnerable (Brasil 2018).

Taxon	English name	Voucher	Migratory status	Sites	% occurrence in 2015–2016	% occurrence in 2017–2018	Record type	Source
<b>Anseriformes Linnaeus, 1758</b>								
<b>Anhimidae Stejneger, 1885</b>								
<i>Chauna torquata</i> (Oken, 1816)	Southern Screamer	WA1961724	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	63	82	BD	1, 2, 5
<b>Anatidae Leach, 1820</b>								
<i>Dendrocygna viduata</i> (Linnaeus, 1766)	White-faced Whistling-Duck	WA2335098	—	S6, T1, T3, T5	4	8	BD	1, 2, 5
<i>Dendrocygna autumnalis</i> (Linnaeus, 1758)	Black-bellied Whistling-Duck	WA2347021	—	T1, T2, T3, T4, T5	8	26	BD	1, 2, 5
<i>Cairina moschata</i> (Linnaeus, 1758)	Muscovy Duck	WA2698723	—	T1, T3, T4, T5	6	10	BD	1, 2, 5
[ <i>Amazonetta brasiliensis</i> (Gmelin, 1789)]	Brazilian Teal		—	T3	—	2	IO	1, 2, 5
<b>Galliformes Linnaeus, 1758</b>								
<b>Cracidae Rafinesque, 1815</b>								
[ <i>Penelope ochrogaster</i> Pelzeln, 1870] VU*	Chestnut-bellied Guan		—		—	—	IO	5
<i>Aburria cumanensis</i> (Jacquin, 1784)	Blue-throated Piping-Guan	WA2698724	—	S1, S2, S4, T1, T2, T3, T4	19	16	BD	1
<i>Aburria cujubi</i> (Pelzeln, 1858)	Red-throated Piping-Guan	WA2028639	—	S1, S2, T1, T2, T3, T4	4	26	BD	1, 2, 5
<i>Ortalis canicollis</i> (Wagler, 1830)	Chaco Chachalaca	WA2080183	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	85	82	BD	1, 2, 5
<i>Crax fasciolata</i> Spix, 1825VU	Bare-faced Curassow	WA2080188	—	S1, S2, S4, T1, T2, T3, T4	13	62	BD	1, 2, 5
<b>Podicipediformes Fürbringer, 1888</b>								
<b>Podicipedidae Bonaparte, 1831</b>								
[ <i>Tachybaptus dominicus</i> (Linnaeus, 1766)]	Least Grebe		—		—	—	IO	2
[ <i>Podilymbus podiceps</i> (Linnaeus, 1758)]	Pied-billed Grebe		—		—	—	IO	5
<b>Ciconiiformes Bonaparte, 1854</b>								
<b>Ciconiidae Sundevall, 1836</b>								
<i>Ciconia maguari</i> (Gmelin, 1789)	Maguari Stork	WA2346180	—	T5	IN	2	BD	1, 5
<i>Jabiru mycteria</i> (Lichtenstein, 1819)	Jabiru	WA2346422	—	S6, T2, T3, T4, T5	2	12	BD	1, 2, 5
<i>Mycteria americana</i> Linnaeus, 1758	Wood Stork	WA3363870	—	T3	6	4	BD	1, 2, 5
<b>Suliformes Sharpe, 1891</b>								
<b>Phalacrocoracidae Reichenbach, 1849</b>								
<i>Nannopterum brasilianus</i> (Gmelin, 1789)	Neotropic Cormorant	WA2706121	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	60	84	BD	1, 2, 5
<b>Anhingidae Reichenbach, 1849</b>								
<i>Anhinga anhinga</i> (Linnaeus, 1766)	Anhinga	WA1986843	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	63	66	BD	1, 2, 5
<b>Pelecaniformes Sharpe, 1891</b>								
<b>Ardeidae Leach, 1820</b>								
<i>Tigrisoma lineatum</i> (Boddaert, 1783)	Rufescent Tiger-Heron	WA2026526	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	65	82	BD	1, 2, 5
<i>Cochlearius cochlearius</i> (Linnaeus, 1766)	Boat-billed Heron	WA2901405	—	S2, S3, S4, T2, T3, T4	17	12	BD	1, 5
<i>Ixobrychus exilis</i> (Gmelin, 1789)	Least Bittern	WA1986845	—	S3, S5, T1, T2, T3, T4, T5	4	72	BD	1, 5
<i>Nycticorax nycticorax</i> (Linnaeus, 1758)	Black-crowned Night-Heron	WA2696526	—	T2, T3, T4, T5	—	12	BD	1, 5
<i>Butorides striata</i> (Linnaeus, 1758)	Striated Heron	WA2031279	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	52	100	BD	1, 5
<i>Bubulcus ibis</i> (Linnaeus, 1758)	Cattle Egret	WA2901507	—	S3, S6, T1, T3	8	14	BD	1, 2
<i>Ardea cocoi</i> Linnaeus, 1766	Cocoi Heron	WA2706120	—	S1, S2, S4, S5, S6, T1, T2, T3, T4, T5	35	68	BD	1, 2, 5
<i>Ardea alba</i> Linnaeus, 1758	Great Egret	WA2102176	—	S5, S6, T1, T2, T3, T4, T5	13	63	BD	1, 2, 5
<i>Pilherodius pileatus</i> (Boddaert, 1783)	Capped Heron	WA2715200	—		IN	IN	BD	1, 2, 5
<i>Egretta thula</i> (Molina, 1782)	Snowy Egret	WA3358045	—	T1, T2, T3, T4, T5	19	30	BD	1, 2, 5
<i>Egretta caerulea</i> (Linnaeus, 1758)	Little Blue Heron	WA2346433	—	S6	2	IN	BD	1
<b>Threskiornithidae Poche, 1904</b>								
<i>Mesembrinibis cayennensis</i> (Gmelin, 1789)	Green Ibis	WA2698730	—	S1, S2, S3, S6, T1, T3	21	6	BD	1, 2, 5
<i>Phimosus infuscatus</i> (Lichtenstein, 1823)	Bare-faced Ibis	WA2698725	—	T1, T2	IN	4	BD	1, 2, 5
<i>Theristicus caerulescens</i> (Vieillot, 1817)	Plumbeous Ibis	WA3358044	—	S4	2	IN	BD	1, 5
[ <i>Theristicus caudatus</i> (Boddaert, 1783)]	Buff-necked Ibis		—	S6	2	IN	IO	1, 2, 5
<i>Platalea ajaja</i> Linnaeus, 1758	Roseate Spoonbill	WA2346434	MPR	S6, T1	6	2	BD	1, 2, 5

Taxon	English name	Voucher	Migratory status	Sites	% occurrence in 2015–2016	% occurrence in 2017–2018	Record type	Source
<b>Cathartiformes Seebohm, 1890</b>								
<b>Cathartidae Lafresnaye, 1839</b>								
<i>Cathartes aura</i> (Linnaeus, 1758)	Turkey Vulture	WA3363006	—	T1, T2, T3, T4, T5	28	36	BD	1, 2, 5
<i>Cathartes burrovianus</i> Cassin, 1845	Lesser Yellow-headed Vulture	WA2102175	—	S4, T1, T2, T3, T4, T5	4	50	BD	1, 2, 5
<i>Coragyps atratus</i> (Bechstein, 1793)	Black Vulture	WA2352546	—	S1, S2, S3, S4, S6, T1, T2, T3, T5	19	42	BD	1, 2, 5
<b>Accipitriformes Bonaparte, 1831</b>								
<b>Pandionidae Bonaparte, 1854</b>								
<i>Pandion haliaetus</i> (Linnaeus, 1758)	Osprey	WA2346063	MGT	T1, T5	IN	10	BD	1
<b>Accipitridae Vigors, 1824</b>								
<i>Leptodon cayanensis</i> (Latham, 1790)	Gray-headed Kite	WA2697657	—	T1	—	2	BD	1
<i>Ictinia mississippiensis</i> (Wilson, 1811)	Mississippi Kite	WA2346791	MGT	S1	3	—	BD	1
<i>Ictinia plumbea</i> (Gmelin, 1788)	Plumbeous Kite	WA3363778	MPR	—	—	IN	BD	1, 5
<i>Busarellus nigricollis</i> (Latham, 1790)	Black-collared Hawk	WA2696527	—	S1, S4, T1, T2, T3, T4, T5	6	42	BD	1, 2, 5
<i>Rostrhamus sociabilis</i> (Vieillot, 1817)	Snail Kite	WA2697716	MPR	S1, S4, T1, T2, T3, T4, T5	6	78	BD	1, 2, 5
<i>Geranospiza caerulescens</i> (Vieillot, 1817)	Crane Hawk	WA3357993	—	T1, T2	19	4	BD	1, 2
<i>Heterospizias meridionalis</i> (Latham, 1790)	Savanna Hawk	WA2346836	—	S3, S6, T1, T2, T3, T4, T5	8	10	BD	1, 2, 5
<i>Urubitinga urubitinga</i> (Gmelin, 1788)	Great Black Hawk	WA2335087	—	S1, S2, S3, S4, T1, T2, T3, T4, T5	15	36	BD	1, 2, 5
<i>Rupornis magnirostris</i> (Gmelin, 1788)	Roadside Hawk	WA2697661	—	S4, T1, T2, T3, T4, T5	4	28	BD	1, 2, 5
<i>Geranoaetus albicaudatus</i> (Vieillot, 1816)	White-tailed Hawk	WA3126426	—	—	—	IN	BD	1, 5
<b>Eurypygiiformes Fürbringer, 1888</b>								
<b>Eurypygidae Selby, 1840</b>								
<i>[Eurypyga helias</i> (Pallas, 1781)]	Sunbittern	—	—	—	—	IN	IO	1
<b>Gruiiformes Bonaparte, 1854</b>								
<b>Aramidae Bonaparte, 1852</b>								
<i>Aramus guarauna</i> (Linnaeus, 1766)	Limpkin	WA2028644	—	S2, S4, S6, T1, T2, T3, T4, T5	15	60	BD	1, 2, 5
<b>Rallidae Rafinesque, 1815</b>								
<i>Aramides cajaneus</i> (Statius Müller, 1776)	Gray-necked Wood-Rail	WA2715199	—	S4, T1, T4	2	6	BD	1, 2, 5
<i>[Mustelirallus albicollis</i> (Vieillot, 1819)]	Ash-throated Crake	—	—	S4, T5	14	6	IO	1, 5
<i>[Pardirallus nigricans</i> (Vieillot, 1819)]	Blackish Rail	—	—	—	—	—	IO	5
<i>Porphyrio martinicus</i> (Linnaeus, 1766)	Purple Gallinule	WA2346181	MPR	T5	IN	2	BD	1, 5
<i>Porphyrio flavirostris</i> (Gmelin, 1789)	Azure Gallinule	WA1952134	ND	S5, S6, T4, T5	13	10	BD	1, 5
<b>Heliornithidae Gray, 1840</b>								
<i>Heliornis fulica</i> (Boddaert, 1783)	Sungrebe	WA1961721	—	S1, S2, S3, S4, S5, T1, T2, T3	27	16	BD	1, 2, 5
<b>Charadriiformes Huxley, 1867</b>								
<b>Charadriidae Leach, 1820</b>								
<i>Vanellus cayanus</i> (Latham, 1790)	Pied Lapwing	WA2698726	—	S6	4	IN	BD	1, 2, 5
<i>Vanellus chilensis</i> (Molina, 1782)	Southern Lapwing	WA2698732	—	S5, S6, T2, T4, T5	6	12	BD	1, 2, 5
<i>Charadrius collaris</i> Vieillot, 1818	Collared Plover	WA2390287	—	S6	4	IN	BD	1, 2
<b>Recurvirostridae Bonaparte, 1831</b>								
<i>Himantopus melanurus</i> Vieillot, 1817	White-backed Stilt	WA2346395	—	S6, T5	2	6	BD	1, 5
<b>Scolopacidae Rafinesque, 1815</b>								
<i>Actitis macularia</i> (Linnaeus, 1766)	Spotted Sandpiper	WA2346027	MGT	S1	2	—	BD	1, 2
<i>Tringa solitaria</i> Wilson, 1813	Solitary Sandpiper	WA2363200	MGT	S6	2	—	BD	1, 2, 5
<i>Tringa flavipes</i> (Gmelin, 1789)	Lesser Yellowlegs	WA2363199	MGT	S6	6	—	BD	1
<i>Calidris fuscicollis</i> (Vieillot, 1819)	White-rumped Sandpiper	WA2901496	MGT	S6	2	IN	BD	1
<i>Calidris melanotos</i> (Vieillot, 1819)	Pectoral Sandpiper	WA2704505	MGT	S6	2	—	BD	1
<i>Phalaropus tricolor</i> (Vieillot, 1819)	Wilson's Phalarope	WA2704503	MGT	S6	2	—	BD	1
<b>Jacanidae Chenu &amp; Des Murs, 1854</b>								
<i>Jacana jacana</i> (Linnaeus, 1766)	Wattled Jacana	WA2028626	—	S5, S6, T1, T2, T3, T4, T5	23	76	BD	1, 2, 5
<b>Laridae Rafinesque, 1815</b>								
<i>Leucophaeus pipixcan</i> (Wagler, 1831)	Franklin's Gull	—	VAG	—	—	—	BD	3
<b>Sternidae Vigors, 1825</b>								
<i>Sternula superciliosa</i> (Vieillot, 1819)	Yellow-billed Tern	WA1953400	—	S6, T2, T3, T5	4	26	BD	1, 2, 5
<i>Phaetusa simplex</i> (Gmelin, 1789)	Large-billed Tern	WA2696528	—	S6, T1, T2, T3, T4, T5	2	64	BD	1, 2, 5
<i>Chlidonias niger</i> (Linnaeus, 1758)	Black Tern	—	MGT	T2	—	2	BD	1
<i>[Sterna hirundo</i> Linnaeus, 1758]	Common Tern	—	MGT	—	—	—	BD	5
<b>Rynchopidae Bonaparte, 1838</b>								
<i>Rynchops niger</i> Linnaeus, 1758	Black Skimmer	WA2698728	MPR	S6, T5	2	8	BD	1, 5

Taxon	English name	Voucher	Migratory status	Sites	% occurrence in 2015–2016	% occurrence in 2017–2018	Record type	Source
<b>Columbiformes Latham, 1790</b>								
<b>Columbidae Leach, 1820</b>								
[ <i>Columbina minuta</i> (Linnaeus, 1766)]	Plain-breasted Ground-Dove		—		IN	IN	IO	1
[ <i>Columbina talpacoti</i> (Temminck, 1810)]	Ruddy Ground-Dove	WA2698731	—	S6	2	IN	BD	1, 2, 5
[ <i>Columbina squammata</i> (Lesson, 1831)]	Scaled Dove		—		—	—	IO	5
[ <i>Columbina picui</i> (Temminck, 1813)]	Picui Ground-Dove	WA3363864	—		IN	IN	BD	1, 2, 5
[ <i>Claravis pretiosa</i> (Ferrari-Perez, 1886)]	Blue Ground-Dove		—	S1, S1	2	2	IO	1, 5
[ <i>Patagioenas picazuro</i> (Temminck, 1813)]	Picazuro Pigeon		—	S3, S4, T1, T2, T3, T4, T5	8	40	IO	1, 2, 5
[ <i>Patagioenas cayennensis</i> (Bonaterre, 1792)]	Pale-vented Pigeon	WA2901501	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	65	82	BD	1, 2, 5
[ <i>Leptotila verreauxi</i> Bonaparte, 1855]	White-tipped Dove	WA3363957	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	69	86	BD	1, 2, 5
[ <i>Leptotila rufaxilla</i> (Richard & Bernard, 1792)]	Gray-fronted Dove		—	S3	14	—	IO	1, 5
<b>Cuculiformes Wagler, 1830</b>								
<b>Cuculidae Leach, 1820</b>								
[ <i>Coccyca minuta</i> (Vieillot, 1817)]	Little Cuckoo	WA1952141	—	S2, S4, S5, T1, T2, T3, T4, T5	17	24	BD	1, 2, 5
[ <i>Playa cayana</i> (Linnaeus, 1766)]	Squirrel Cuckoo		—	S3, S4, T1, T2, T3, T4	13	18	IO	1, 5
[ <i>Crotophaga major</i> Gmelin, 1788]	Greater Ani	WA2028628	—	S1, S2, S3, S4, T1, T2, T3, T4, T5	50	54	BD	1, 2, 5
[ <i>Crotophaga ani</i> Linnaeus, 1758]	Smooth-billed Ani	WA2028634	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	73	86	BD	1, 2, 5
[ <i>Guira guira</i> (Gmelin, 1788)]	Guira Cuckoo	WA3363958	—	T5	IN	10	BD	1, 5
[ <i>Tapera naevia</i> (Linnaeus, 1766)]	Striped Cuckoo		—	S1, S5, S6, T1, T2, T3, T4, T5	17	30	IO	1, 5
<b>Strigiformes Wagler, 1830</b>								
<b>Tytonidae Mathews, 1912</b>								
[ <i>Tyto furcata</i> (Temminck, 1827)]	American Barn Owl	WA2972378	—		IN	IN	BD	1, 2
<b>Strigidae Leach, 1820</b>								
[ <i>Megascops choliba</i> (Vieillot, 1817)]	Tropical Screech-Owl		—		—	—	IO	5
[ <i>Bubo virginianus</i> (Gmelin, 1788)]	Great Horned Owl		—	S3	2	IN	IO	1, 5
[ <i>Glaucidium brasilianum</i> (Gmelin, 1788)]	Ferruginous Pygmy-Owl		—	S1, S3, T1, T3, T4	6	6	IO	1, 5
<b>Nyctibiiformes Yuri et al., 2013</b>								
<b>Nyctibiidae Chenu &amp; Des Murs, 1851</b>								
[ <i>Nyctibius griseus</i> (Gmelin, 1789)]	Common Potoo	WA3363846	—	S4	2	—	BD	1, 5
<b>Caprimulgiformes Ridgway, 1881</b>								
<b>Caprimulgidae Vigers, 1825</b>								
[ <i>Nyctiprogne leucopyga</i> (Spix, 1825)]	Band-tailed Nighthawk		—	T3, S1	IN	4	IO	1
[ <i>Nyctidromus albicollis</i> (Gmelin, 1789)]	Common Pauraque	WA2715203	—	S3, T3, T4	6	10	BD	1, 2, 5
[ <i>Hydropsalis parvula</i> (Gould, 1837)]	Little Nightjar		MPR		—	—	IO	2, 5
[ <i>Podager nacunda</i> (Vieillot, 1817)]	Nacunda Nighthawk		MPR		—	IN	IO	1, 2, 5
<b>Apodiformes Peters, 1940</b>								
<b>Apodidae Olphe-Galliard, 1887</b>								
[ <i>Streptoprocne zonaris</i> (Shaw, 1796)]	White-collared Swift	WA3363751	—	T3	—	2	BD	1
<b>Trochilidae Vigers, 1825</b>								
[ <i>Glaucis hirsutus</i> (Gmelin, 1788)]	Rufous-breasted Hermit		—		—	—	IO	5
[ <i>Phaethornis ruber</i> (Linnaeus, 1758)]	Reddish Hermit		—		—	—	IO	5
[ <i>Phaethornis subochraceus</i> Todd, 1915]	Buff-bellied Hermit	WA2336756	—	S3, S4, T1, T2, T3, T4	6	14	BD	1, 5
[ <i>Phaethornis pretrei</i> (Lesson & Delattre, 1839)]	Planalto Hermit		—	S3	3	—	IO	1, 2, 5
[ <i>Chlorostilbon lucidus</i> (Shaw, 1812)]	Glittering-bellied Emerald		—		—	—	IO	2, 5
[ <i>Thalurania furcata</i> (Gmelin, 1788)]	Fork-tailed Woodnymph	WA3357991	—	T3	—	4	BD	1, 5
[ <i>Hylocharis cyanus</i> (Vieillot, 1818)]	White-chinned Sapphire		—		—	—	IO	2
[ <i>Hylocharis chrysura</i> (Shaw, 1812)]	Gilded Hummingbird	WA2336899	—	S1, S4, T1, T2, T3, T4	4	14	BD	1, 2, 5
[ <i>Polytmus gainumbi</i> (Pallas, 1764)]	White-tailed Goldenthrout	WA2697678	—	T5, S5	8	4	BD	1, 5
[ <i>Amazilia versicolor</i> (Vieillot, 1818)]	Versicolored Emerald		—		—	—	IO	2, 5
[ <i>Amazilia fimbriata</i> (Gmelin, 1788)]	Glittering-throated Emerald		—		—	—	IO	2, 5
<b>Trogoniformes A. O. U., 1886</b>								
<b>Trogonidae Lesson, 1828</b>								
[ <i>Trogon curucui</i> Linnaeus, 1766]	Blue-crowned Trogon	WA1986817	—	S1, S2, S3, S4, T1, T2, T3, T4	33	34	BD	1, 2, 5
<b>Coraciiformes Forbes, 1844</b>								
<b>Alcedinidae Rafinesque, 1815</b>								
[ <i>Megaceryle torquata</i> (Linnaeus, 1766)]	Ringed Kingfisher	WA2901877	—	S1, S2, S3, S4, S5, T1, T2, T3, T4	23	42	BD	1, 2, 5

Taxon	English name	Voucher	Migratory status	Sites	% occurrence in 2015–2016	% occurrence in 2017–2018	Record type	Source
<i>Chloroceryle amazona</i> (Latham, 1790)	Amazon Kingfisher	WA2697688	—	S1, T1, T2, T3, T4	4	20	BD	1, 2, 5
<i>Chloroceryle aenea</i> (Pallas, 1764)	American Pygmy Kingfisher	WA2390288	—	S3, S6, T3, T4	6	8	BD	1, 5
<i>Chloroceryle americana</i> (Gmelin, 1788)	Green Kingfisher	WA2347015	—	S1, S3, S4, T1, T2, T3, T4	13	42	BD	1, 5
[ <i>Chloroceryle inda</i> (Linnaeus, 1766)]	Green-and-rufous Kingfisher		—	S2, T2	4	2	IO	1, 5
<b>Momotidae Gray, 1840</b>								
[ <i>Momotus momota</i> (Linnaeus, 1766)]	Amazonian Motmot		—		—	—	IO	2
<b>Galbuliformes Fürbringer, 1888</b>								
<b>Galbulidae Vigors, 1825</b>								
<i>Galbula ruficauda</i> Cuvier, 1816	Rufous-tailed Jacamar	WA2080244	—	S1, S2, S3, S4, S6, T1, T2, T3, T4	48	66	BD	1, 2, 5
<b>Bucconidae Horsfield, 1821</b>								
<i>Monasa nigrifrons</i> (Spix, 1824)	Black-fronted Nunbird	WA2704528	—	S1, S2, S3, S4, T1, T2, T3, T4	29	30	BD	1, 2, 5
<b>Piciformes Meyer &amp; Wolf, 1810</b>								
<b>Ramphastidae Vigors, 1825</b>								
<i>Ramphastos toco</i> Statius Müller, 1776	Toco Toucan	WA2901404	—	S1, S2, S3, T1, T4	8	6	BD	1, 2, 5
<i>Pteroglossus castanotis</i> Gould, 1834	Chestnut-eared Aracari	WA3357747	—	S3, S4, T2, T3	4	6	BD	1
<b>Picidae Leach, 1820</b>								
<i>Picumnus albosquamatus</i> d'Orbigny, 1840	White-wedged Piculet	WA3357982	—	T1, T2, T3, T4	—	12	BD	1, 5
[ <i>Melanerpes cruentatus</i> (Boddaert, 1783)]	Yellow-tufted Woodpecker		—		—	—	IO	5
<i>Veniliornis passerinus</i> (Linnaeus, 1766)	Little Woodpecker	WA2901406	—	S1, S2, S3, S4, T1, T2, T3, T4	17	20	BD	1, 2, 5
[ <i>Piculus chrysochloros</i> (Vieillot, 1818)]	Golden-green Woodpecker		—	S1, S3, S4, T1, T2	10	4	IO	1, 5
[ <i>Colaptes melanochloros</i> (Gmelin, 1788)]	Green-barred Woodpecker		—	S1	2	IN	IO	1, 5
<i>Ceuleus lugubris</i> (Malherbe, 1851)	Pale-crested Woodpecker	WA2901876	—	S1, S4, T1, T2, T4	6	10	BD	1, 2, 5
<i>Ceuleus flavus</i> (Statius Müller, 1776)	Cream-colored Woodpecker	WA2901875	—	S4, T1, T2, T3, T4	2	20	BD	1, 5
[ <i>Campephilus melanoleucos</i> (Gmelin, 1788)]	Crimson-crested Woodpecker		—	S1, S2, T1, T2	10	6	IO	1, 2, 5
<b>Falconiformes Bonaparte, 1831</b>								
<b>Falconidae Leach, 1820</b>								
<i>Caracara plancus</i> (Miller, 1777)	Southern Caracara	WA1999800	—	S1, S2, S4, S5, S6, T2, T3, T4, T5	19	36	BD	1, 2, 5
[ <i>Milvago chimachima</i> (Vieillot, 1816)]	Yellow-headed Caracara		—		—	—	IO	5
[ <i>Herpetotheres cachinnans</i> (Linnaeus, 1758)]	Laughing Falcon		—	S4, T3	2	4	IO	1, 5
[ <i>Falco sparverius</i> Linnaeus, 1758]	American Kestrel		—		—	—	IO	5
<i>Falco rufigularis</i> Daudin, 1800	Bat Falcon	WA2706122	—	S4	2	—	BD	1
<b>Psittaciformes Wagler, 1830</b>								
<b>Psittacidae Rafinesque, 1815</b>								
<i>Ara ararauna</i> (Linnaeus, 1758)	Blue-and-yellow Macaw	WA2901436	—	T3, S1	—	6	BD	1
<i>Primolius auricollis</i> (Cassin, 1853)	Yellow-collared Macaw	WA3363933	—	S2, T1, T3, S3	2	10	BD	1, 5
[ <i>Diopsittaca nobilis</i> (Linnaeus, 1758)]	Red-shouldered Macaw		—	S1, S2, S3, S4, T2, T4	8	12	IO	1, 2, 5
<i>Thectocercus acuticaudatus</i> (Vieillot, 1818)	Blue-crowned Parakeet	WA2697662	—	T1, T2, T3, T4	—	24	BD	1
<i>Psittacara leucophthalmus</i> (Statius Müller, 1776)	White-eyed Parakeet	WA2033574	—	T1, T2, T3, T4	14	12	BD	1, 2, 5
[ <i>Aratinga nenday</i> (Vieillot, 1823)]	Nanday Parakeet		—		—	IN	IO	1, 2, 5
[ <i>Eupsittula aurea</i> (Gmelin, 1788)]	Peach-fronted Parakeet		—		—	4	IO	2, 5
<i>Myiopsitta monachus</i> (Boddaert, 1783)	Monk Parakeet	WA2031283	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	54	96	BD	1, 2, 5
[ <i>Forpus xanthopterygius</i> (Spix, 1824)]	Blue-winged Parrotlet		—		—	—	IO	2, 5
<i>Brotogeris chiriri</i> (Vieillot, 1818)	Yellow-chevroned Parakeet	WA3357746	—	S2, S3, T1, T2, T3, T4, T5	6	36	BD	1, 2, 5
[ <i>Pionus menstruus</i> (Linnaeus, 1766)]	Blue-headed Parrot		—		—	IN	IO	1
<i>Pionus maximiliani</i> (Kuhl, 1820)	Scaly-headed Parrot	WA2141194	—	S4	6	IN	BD	1, 2, 5
[ <i>Amazona amazonica</i> (Linnaeus, 1766)]	Orange-winged Parrot		—	S1	6	IN	IO	1, 2, 5
<i>Amazona aestiva</i> (Linnaeus, 1758)	Turquoise-fronted Parrot	WA2901499	—	S1, S2, S3, S4, S6, T1, T2, T3, T4, T5	44	70	BD	1, 2, 5
<b>Passeriformes Linnaeus, 1758</b>								
<b>Thamnophilidae Swainson, 1824</b>								
[ <i>Herpilochmus longirostris</i> Pelzeln, 1868]	Large-billed Antwren		—		—	—	IO	5
[ <i>Thamnophilus doliatus</i> (Linnaeus, 1764)]	Barred Antshrike		—	S2, T2, T3, T4	2	12	IO	1, 5
[ <i>Thamnophilus sticturus</i> Pelzeln, 1868]	Bolivian Slaty-Antshrike		—		—	—	IO	2
[ <i>Thamnophilus caerulescens</i> Vieillot, 1816]	Variable Antshrike		—		—	—	IO	5
<i>Taraba major</i> (Vieillot, 1816)	Great Antshrike	WA2080191	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	83	96	BD	1, 2, 5
<i>Hypocnemoides maculicauda</i> (Pelzeln, 1868)	Band-tailed Antbird	WA3041345	—	S3, T1, T2, T3, T4	2	10	BD	1, 5

Taxon	English name	Voucher	Migratory status	Sites	% occurrence in 2015–2016	% occurrence in 2017–2018	Record type	Source
[ <i>Pyriglena leuconota</i> (Spix, 1824)]	White-backed Fire-eye		—		—	—	IO	5
<i>Cercomacra melanaria</i> (Ménétrières, 1835)	Mato Grosso Antbird	WA2141195	—	S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	60	62	BD	1, 2, 5
<b>Dendrocolaptidae Gray, 1840</b>								
[ <i>Sittasomus griseicapillus</i> (Vieillot, 1818)]	Olivaceous Woodcreeper		—	T2, S1	—	4	IO	1, 2, 5
<i>Xiphorhynchus guttatoides</i> (Lafresnaye, 1850)	Lafresnaye's Woodcreeper	WA2901880	—	S1, S4, T3	4	2	BD	1
<i>Campylorhamphus trochilirostris</i> (Lichtenstein, 1820)	Red-billed Scythebill	WA2901500	—	S1, S3, T1, T2, T3, T4, T5	6	20	BD	1, 2
<i>Dendroplex picus</i> (Gmelin, 1788)	Straight-billed Woodcreeper	WA3358042	—	S1, S2, T1, T3, T4	6	12	BD	1, 2, 5
[ <i>Lepidocolaptes angustirostris</i> (Vieillot, 1818)]	Narrow-billed Woodcreeper		—		—	—	IO	5
<b>Furnariidae Gray, 1840</b>								
<i>Furnarius leucopus</i> Swainson, 1838	Pale-legged Hornero	WA2698729	—	S1, S2, S3, S4, S6, T1, T2, T3, T4, T5	52	82	BD	1, 2, 5
[ <i>Furnarius rufus</i> (Gmelin, 1788)]	Rufous Hornero		—		—	—	IO	2, 5
<i>Pseudoseisura unirufa</i> (d'Orbigny & Lafresnaye, 1838)	Rufous Cacholote	WA1952114	—	S2, S3, S4, S6, T1, T2, T3, T4	15	14	BD	1, 2, 5
[ <i>Phacellodomus rufifrons</i> (Wied, 1821)]	Rufous-fronted Thornbird		—		—	—	IO	2, 5
<i>Phacellodomus ruber</i> (Vieillot, 1817)	Greater Thornbird	WA2390289	—	S1, S3, S6, T4	6	2	BD	1, 5
<i>Certhiaxis cinnamomeus</i> (Gmelin, 1788)	Yellow-chinned Spinetail	WA2031282	—	S1, S2, S3, S5, S6, T1, T2, T3, T4, T5	42	98	BD	1, 2, 5
<i>Synallaxis hypospodia</i> Sclater, 1874	Cinereous-breasted Spinetail	WA2365310	—	S5, T3, T4, T5	2	20	BD	1
<i>Synallaxis albilora</i> Pelzeln, 1856	White-lored Spinetail	WA2905329	—	S1, S2, S4, T1, T2, T3, T4, T5	17	66	BD	1, 2, 5
<i>Cranioleuca vulpina</i> (Pelzeln, 1856)	Rusty-backed Spinetail	WA2141187	—	S1, S2, S3, S4, T1, T3, T4	21	10	BD	1, 2
<b>Tityridae Gray, 1840</b>								
<i>Pachyrhamphus polychopterus</i> (Vieillot, 1818)	White-winged Becard	WA3363852	MPR	S2	2	—	BD	1, 5
[ <i>Xenopsaris albinucha</i> (Burmeister, 1869)]	White-naped Xenopsaris		—		—	—	IO	5
Rhynchocyclidae Berlepsch, 1907								
[ <i>Tolmomyias sulphurescens</i> (Spix, 1825)]	Yellow-olive Flycatcher		—	T2, T3	—	4	IO	1
<i>Todirostrum cinereum</i> (Linnaeus, 1766)	Common Tody-Flycatcher	WA2035462	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4	31	52	BD	1, 2, 5
<i>Poecilotriccus latirostris</i> (Pelzeln, 1868)	Rusty-fronted Tody-Flycatcher	WA2697679	—	S3, S4, T1, T2, T3, T4	6	18	BD	1, 2, 5
[ <i>Hemitriccus margaritaceiventer</i> (d'Orbigny & Lafresnaye, 1837)]	Pearly-vented Tody-tyrant		—		—	—	IO	5
<b>Tyrannidae Vigors, 1825</b>								
[ <i>Inezia inornata</i> (Salvadori, 1897)]	Plain Tyrannulet		MGT	T2, T3	—	4	IO	1, 5
[ <i>Euscarthmus meloryphus</i> Wied, 1831]	Tawny-crowned Pygmy-Tyrant		—		—	—	IO	5
[ <i>Camptostoma obsoletum</i> (Temminck, 1824)]	Southern Beardless-Tyrannulet		—	S4, S1	2	2	IO	1, 2, 5
[ <i>Elaenia flavogaster</i> (Thunberg, 1822)]	Yellow-bellied Elaenia		—		—	—	IO	2, 5
<i>Elaenia spectabilis</i> Pelzeln, 1868	Large Elaenia	WA2346878	MPR	S1, S2, T4	6	2	BD	1
[ <i>Elaenia cristata</i> Pelzeln, 1868]	Plain-crested Elaenia		—		—	—	IO	5
[ <i>Elaenia chiriquensis</i> Lawrence, 1865]	Lesser Elaenia		MPR		—	—	IO	5
[ <i>Myiopagis gaimardii</i> (d'Orbigny, 1839)]	Forest Elaenia		—		—	—	IO	5
[ <i>Myiopagis caniceps</i> (Swainson, 1835)]	Gray Elaenia		—		—	—	IO	5
[ <i>Serpophaga subcristata</i> (Vieillot, 1817)]	White-crested Tyrannulet		—		—	—	IO	5
[ <i>Attila bolivianus</i> Lafresnaye, 1848]	Dull-capped Attila		—	S3, S4	6	—	IO	1
[ <i>Legatus leucophaius</i> (Vieillot, 1818)]	Piratic Flycatcher		MPR		—	—	IO	5
<i>Myiarchus ferox</i> (Gmelin, 1789)	Short-crested Flycatcher		—	S2, S4, T1, T2, T3, T4	4	14	AD	1, 2, 4
[ <i>Myiarchus tyrannulus</i> (Statius Müller, 1776)]	Brown-crested Flycatcher		—		—	—	IO	5
<i>Casiornis rufus</i> (Vieillot, 1816)	Rufous Casiornis	WA2696556	—	T3	—	2	BD	1, 2
<i>Pitangus sulphuratus</i> (Linnaeus, 1766)	Great Kiskadee	WA2704529	MPR	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4	100	100	BD	1, 2, 5
<i>Philohydor lictor</i> (Lichtenstein, 1823)	Lesser Kiskadee	WA2352668	—	S1, S4, T2, T3	6	20	BD	1, 5
[ <i>Machetornis rixosa</i> (Vieillot, 1819)]	Cattle Tyrant		—		—	—	IO	5
[ <i>Megarynchus pitangua</i> (Linnaeus, 1766)]	Boat-billed Flycatcher		—	S4, T1, T2, T3, T4	6	16	IO	1, 2, 5
<i>Myiozetetes cayanensis</i> (Linnaeus, 1766)	Rusty-margined Flycatcher	WA1986818	—	S1, S2, S3, S4, S6, T1, T2, T3, T4	65	42	BD	1, 2, 5
[ <i>Tyrannus albogularis</i> Burmeister, 1856]	White-throated Kingbird		MPR		—	—	IO	5
<i>Tyrannus melancholicus</i> Vieillot, 1819	Tropical Kingbird	WA1998625	MPR	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	81	92	BD	1, 2, 5



Taxon	English name	Voucher	Migratory status	Sites	% occurrence in 2015–2016	% occurrence in 2017–2018	Record type	Source
[ <i>Tyrannus savana</i> Daudin, 1802]	Fork-tailed Flycatcher		MPR		—	IN	IO	1
[ <i>Tyrannus tyrannus</i> (Linnaeus, 1758)]	Eastern Kingbird		MGT		—	—	IO	5
[ <i>Griseotyrannus aurantioatrocristatus</i> (d'Orbigny & Lafresnaye, 1837)]	Crowned Slaty Flycatcher		MPR		—	—	IO	5
<i>Empidonomus varius</i> (Vieillot, 1818)	Variegated Flycatcher	WA2724800	MPR		—	IN	AD	1, 4, 5
[ <i>Colonia colonus</i> (Vieillot, 1818)]	Long-tailed Tyrant		—		—	—	IO	5
[ <i>Myiophobus fasciatus</i> (Statius Müller, 1776)]	Bran-colored Flycatcher		MPR		—	—	IO	5
[ <i>Sublegatus modestus</i> (Wied, 1831)]	Southern Scrub-Flycatcher		MPR		—	—	IO	2, 5
<i>Pyrocephalus rubinus</i> (Boddaert, 1783)	Vermilion Flycatcher	WA3357752	MPR	S6, T1	2	2	BD	1, 2, 5
<i>Fluvicola albiventer</i> (Spix, 1825)	Black-backed Water-Tyrant	WA2901460	MPR	S1, S5, S6, T1, T3, T4, T5	10	26	BD	1, 2
<i>Arundinicola leucocephala</i> (Linnaeus, 1764)	White-headed Marsh Tyrant	WA2031280	—	S1, S5, S6, T5	21	4	BD	1, 5
[ <i>Gubernetes yetapa</i> (Vieillot, 1818)]	Streamer-tailed Tyrant		—		—	—	IO	5
<i>Cnemotriccus fuscatus</i> (Wied, 1831)	Fuscous Flycatcher	WA3041344	—	S2	—	4	BD	1, 2, 5
[ <i>Knipolegus hudsoni</i> Sclater, 1872]	Hudson's Black-Tyrant		VAG		—	—	IO	2, 5
<i>Satrapa icterophrys</i> (Vieillot, 1818)	Yellow-browed Tyrant	WA2696555	—	T5	—	8	BD	1, 5
[ <i>Xolmis irupero</i> (Vieillot, 1823)]	White Monjita		—		—	—	IO	5
<b>Vireonidae Swainson, 1837</b>								
<i>Cyclarhis gujanensis</i> (Gmelin, 1789)	Rufous-browed Peppershrike	WA2704502	—	S1, S2, S3, S4, T1, T2, T3, T4	46	46	BD	1, 2, 5
<i>Hylophilus pectoralis</i> Sclater, 1866	Ashy-headed Greenlet	WA3363016	—	S3, T2, T3	2	10	BD	1, 2, 5
[ <i>Vireo olivaceus</i> (Linnaeus, 1766)]	Red-eyed Vireo		MGT		—	—	IO	5
<b>Corvidae Leach, 1820</b>								
[ <i>Cyanocorax cyanomelas</i> (Vieillot, 1818)]	Purplish Jay		—	S1, S3	6	—	IO	1
<b>Hirundinidae Rafinesque, 1815</b>								
<i>Stelgidopteryx ruficollis</i> (Vieillot, 1817)	Southern Rough-winged Swallow	WA2697690	MPR	S1, S2, S4, T1, T2, T3, T4, T5	10	24	BD	1, 2, 5
<i>Progne tapera</i> (Vieillot, 1817)	Brown-chested Martin	WA2698727	MPR	T3, T5	31	10	BD	1, 2, 5
[ <i>Progne subis</i> (Linnaeus, 1758)]	Purple Martin		MGT		IN	—	IO	1
[ <i>Progne chalybea</i> (Gmelin, 1789)]	Gray-breasted Martin		MPR		—	8	IO	2, 5
<i>Tachycineta albiventer</i> (Boddaert, 1783)	White-winged Swallow	WA2907502	—	S4, T1, T2, T3, T4, T5	2	16	BD	1, 2, 5
[ <i>Tachycineta leucorrhoa</i> (Vieillot, 1817)]	White-rumped Swallow		—		—	IN	IO	1, 2, 5
<i>Riparia riparia</i> (Linnaeus, 1758)	Bank Swallow	WA3363003	MGT	S1	6	—	BD	1
<i>Hirundo rustica</i> Linnaeus, 1758	Barn Swallow		MGT	S5, T2, T3, T5	13	14	BD	1
<i>Petrochelidon pyrrhonota</i> (Vieillot, 1817)	Cliff Swallow		MPR	S5, T2, T3, T5	17	28	BD	1
<b>Troglodytidae Swainson, 1831</b>								
[ <i>Troglodytes musculus</i> Naumann, 1823]	Southern House Wren		—		—	—	IO	5
<i>Campylorhynchus turdinus</i> (Wied, 1831)	Thrush-like Wren	WA2815039	—	S1, S2, S3, S4, S6, T1, T2, T3, T4, T5	77	78	BD	1, 2, 5
[ <i>Pheugopedius genibarbis</i> (Swainson, 1838)]	Moustached Wren		—	S4, T2, T3, T4	4	10	IO	1, 2, 5
[ <i>Cantorchilus leucotis</i> (Lafresnaye, 1845)]	Buff-breasted Wren		—	S3	2	IN	IO	1, 2, 5
<b>Donacobiidae Aleixo &amp; Pacheco, 2006</b>								
<i>Donacobius atricapilla</i> (Linnaeus, 1766)	Black-capped Donacobius	WA1961723	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	58	100	BD	1, 2, 5
<b>Poliophtidae Baird, 1858</b>								
<i>Poliophtila dumicola</i> (Vieillot, 1817)	Masked Gnatcatcher	WA2905054	—	T1, T3	—	6	BD	1, 2, 5
<b>Turdidae Rafinesque, 1815</b>								
[ <i>Turdus leucomelas</i> Vieillot, 1818]	Pale-breasted Thrush		—	S1	4	—	IO	1, 5
<i>Turdus rufiventris</i> Vieillot, 1818	Rufous-bellied Thrush	WA2704809	—	S1, S3, S4, T2, T4	13	6	BD	1, 2, 5
<i>Turdus amaurochalinus</i> Cabanis, 1850	Creamy-bellied Thrush	WA3357978	MPR	T1, T2, T5	—	8	BD	1, 2, 5
<b>Mimidae Bonaparte, 1853</b>								
[ <i>Mimus saturninus</i> (Lichtenstein, 1823)]	Chalk-browed Mockingbird		—		—	—	IO	2, 5
<b>Motacillidae Horsfield, 1821</b>								
[ <i>Anthus lutescens</i> Pucheran, 1855]	Yellowish Pipit		—	S6	2	—	IO	1
<b>Passerellidae Cabanis &amp; Heine, 1850</b>								
[ <i>Zonotrichia capensis</i> (Statius Müller, 1776)]	Rufous-collared Sparrow		—		—	—	IO	5
<b>Parulidae et al., 1947</b>								
<i>Setophaga pitiayumi</i> (Vieillot, 1817)	Tropical Parula	WA3357779	—	T3	—	2	BD	1, 2, 5
<i>Geothlypis aequinoctialis</i> (Gmelin, 1789)	Masked Yellowthroat	WA2696554	—	T2, T3, T4, T5	—	16	BD	1, 5
[ <i>Myiothlypis flaveola</i> Baird, 1865]	Flavescent Warbler		—		—	—	IO	5
<b>Icteridae Vigors, 1825</b>								
[ <i>Psarocolius decumanus</i> (Pallas, 1769)]	Crested Oropendola		—	T1, T2, T5	—	6	IO	1, 2
<i>Proccacius solitarius</i> (Vieillot, 1816)	Solitary Black Cacique	WA1998619	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	83	98	BD	1, 2, 5

Taxon	English name	Voucher	Migratory status	Sites	% occurrence in 2015–2016	% occurrence in 2017–2018	Record type	Source
<i>Cacicus cela</i> (Linnaeus, 1758)	Yellow-rumped Cacique	WA1999799	—	S1, S2, S3, S4, S6, T1, T2, T3, T4, T5	69	84	BD	1, 2, 5
[ <i>Icterus pyrrhopterus</i> (Vieillot, 1819)]	Variable Oriole	—	—	S2, S3, S4, S6, T1, T2, T3, T4	31	24	IO	1, 2, 5
<i>Icterus croconotus</i> (Wagler, 1829)	Orange-backed Troupial	WA2901878	—	S1, S2, S3, S4, S6, T2, T3, T4	65	78	BD	1, 2, 5
[ <i>Gnorimopsar chopi</i> (Vieillot, 1819)]	Chopi Blackbird	—	—	—	—	—	IO	5
<i>Amblyramphus holosericeus</i> (Scopoli, 1786)	Scarlet-headed Blackbird	WA1961722	—	T3, T4, T5	6	10	BD	1, 5
<i>Agelasticus cyanopus</i> (Vieillot, 1819)	Unicolored Blackbird	WA2080187	—	S1, S2, S3, S5, S6	56	96	AD	1, 2, 4, 5
[ <i>Agelaioides badius</i> (Vieillot, 1819)]	Grayish Baywing	—	—	—	—	—	IO	2, 5
<i>Molothrus oryzivorus</i> (Gmelin, 1788)	Giant Cowbird	WA3363866	—	—	IN	—	BD	1, 2, 5
[ <i>Molothrus bonariensis</i> (Gmelin, 1789)]	Shiny Cowbird	—	—	—	IN	—	IO	1, 2, 5
[ <i>Sturnella supercilialis</i> (Bonaparte, 1850)]	White-browed Meadowlark	—	—	—	—	—	IO	2
<b>Thraupidae Cabanis, 1847</b>								
[ <i>Paroaria coronata</i> (Miller, 1776)]	Red-crested Cardinal	—	—	—	—	—	IO	5
<i>Paroaria capitata</i> (d'Orbigny & Lafresnaye, 1837)	Yellow-billed Cardinal	WA1952236	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	79	94	AD	1, 2, 4, 5
[ <i>Tangara sayaca</i> (Linnaeus, 1766)]	Sayaca Tanager	—	—	S2, T1, T2, T3, T4	2	30	IO	1, 2, 5
<i>Tangara palmarum</i> (Wied, 1821)	Palm Tanager	WA2706124	—	S3, T1	4	2	BD	1, 5
[ <i>Nemosia pileata</i> (Boddaert, 1783)]	Hooded Tanager	—	—	—	—	—	IO	5
[ <i>Conirostrum speciosum</i> (Temminck, 1824)]	Chestnut-vented Conebill	—	—	—	—	—	IO	2, 5
[ <i>Sicalis flaveola</i> (Linnaeus, 1766)]	Saffron Finch	—	—	—	—	—	IO	5
[ <i>Volatinia jacarina</i> (Linnaeus, 1766)]	Blue-black Grassquit	—	—	—	—	IN	IO	1, 2, 5
[ <i>Eucometis penicillata</i> (Spix, 1825)]	Gray-headed Tanager	—	—	—	—	—	IO	5
<i>Coryphospingus cucullatus</i> (Statius Müller, 1776)	Red-crested Finch	—	—	—	—	—	AD	2, 4, 5
[ <i>Tachyphonus rufus</i> (Boddaert, 1783)]	White-lined Tanager	—	—	—	—	—	IO	2, 5
<i>Ramphocelus carbo</i> (Pallas, 1764)	Silver-beaked Tanager	WA2704530	—	S1, S2, S3, S4, S5, S6, T2, T3, T4	67	72	BD	1, 2, 5
[ <i>Dacnis cayana</i> (Linnaeus, 1766)]	Blue Dacnis	—	—	—	—	—	IO	5
[ <i>Sporophila lineola</i> (Linnaeus, 1758)]	Lined Seedeater	—	MPR	T5	IN	6	IO	1, 5
<i>Sporophila collaris</i> (Boddaert, 1783)	Rusty-collared Seedeater	WA1952061	—	S1, S5, S6, T2, T3, T4, T5	29	32	BD	1, 5
[ <i>Sporophila nigricollis</i> (Vieillot, 1823)]	Yellow-bellied Seedeater	—	—	—	—	—	IO	5
<i>Sporophila caerulescens</i> (Vieillot, 1823)	Double-collared Seedeater	WA2901879	MPR	S6, T2, T3, T4, T5	4	18	BD	1, 5
<i>Sporophila leucoptera</i> (Vieillot, 1817)	White-bellied Seedeater	WA2697745	—	S3, S5, T4, T5	8	4	BD	1, 5
[ <i>Sporophila nigrorufa</i> (d'Orbigny & Lafresnaye, 1837)]	Black-and-tawny Seedeater	—	—	—	—	—	IO	5
[ <i>Sporophila bouvreuil</i> (Statius Müller, 1776)]	Copper Seedeater	—	MPR	—	—	—	IO	5
[ <i>Sporophila hypoxantha</i> Cabanis, 1851]	Tawny-bellied Seedeater	—	MPR	—	—	IN	IO	1, 5
[ <i>Sporophila angolensis</i> (Linnaeus, 1766)]	Chestnut-bellied Seed-Finch	—	—	—	—	—	IO	5
<i>Saltator coerulescens</i> Vieillot, 1817	Grayish Saltator	WA2697740	—	S1, S2, S3, S4, S5, S6, T1, T2, T3, T4, T5	77	90	AD	1, 2, 4, 5
[ <i>Saltator similis</i> d'Orbigny & Lafresnaye, 1837]	Green-winged Saltator	—	—	S2, S3, T3, T5	6	6	IO	1, 5
[ <i>Thlypsopsis sordida</i> (d'Orbigny & Lafresnaye, 1837)]	Orange-headed Tanager	—	—	—	IN	IN	IO	1, 2, 5
<b>Fringillidae Leach, 1820</b>								
[ <i>Euphonia chlorotica</i> (Linnaeus, 1766)]	Purple-throated Euphonia	—	—	S2, S3, T3	4	2	IO	1, 5
[ <i>Passer domesticus</i> (Linnaeus, 1758)]	House Sparrow	—	—	—	—	—	IO	5

were Tyrannidae (18 species), followed by Thraupidae (15), Psittacidae (12), and Ardeidae (11). Non-passerine families were more representative (67%) than passerine families (33%). Of the 34 migrant species, 13 were migratory and 21 were partially migratory. Only *Crax fasciolata* Spix, 1825 has been globally classified as Vulnerable (IUCN 2019).

*Pitangus sulphuratus* (Linnaeus, 1766) was recorded in all samples. *Butorides striata* (Linnaeus, 1758) and *Donacobius atricapilla* (Linnaeus, 1766) were recorded in all samples of the 2017–2018 cycle. Other species with a high frequency of occurrence and which were very

common in EET were *Nannopterum brasilianus* (Gmelin, 1789), *Taraba major* (Vieillot, 1816), *Agelasticus cyanopus* (Vieillot, 1819), *Procacicus solitarius* (Vieillot, 1816), and *Saltator coerulescens* Vieillot, 1817, as well as others in Table 2.

For the less frequent category, we highlight species such as *Platalea ajaja* Linnaeus, 1758 (Fig. 2A) and *Rynchops niger* Linnaeus, 1758 (Fig. 2B), which are partially migratory, and *Phalaropus tricolor* (Vieillot, 1819) and *Hirundo rustica* Linnaeus, 1758, which are migratory. Also among the lower frequency species are *Ciconia maguari* (Gmelin, 1789) (Fig. 2C) and *Amblyramphus*



**Figure 2.** A. *Platalea ajaja* Linnaeus, 1758. B. *Rynchops niger* Linnaeus, 1758. C. *Ciconia maguari* (Gmelin, 1789). D. *Amblyramphus holosericeus* (Scopoli, 1786). E. *Actitis macularius* (Linnaeus, 1766). F. *Ixobrychus exilis* (Gmelin, 1789). (Photos: BDV, except for *Actitis macularius*, photographed by AVBF).

*holosericeus* (Scopoli, 1786) (Fig. 2D), which have been recorded specifically in flooded grassland or floating meadows.

**NEW RECORD FOR THE PANTANAL WETLAND AND THE CENTRAL REGION OF BRAZIL**

Order Charadriiformes  
 Family Sternidae

***Chlidonias niger* (Linnaeus, 1758)**


Figure 3

**Materials examined.** Tables 1, 2.

**Identification.** In April 2018, we saw two non-breeding individuals of *C. niger* flying on the river (16°52'S, 057°24'W) together with *Phaetusa simplex* (Gmelin, 1789) and *Sternula superciljaris* (Vieillot, 1819). The



**Figure 3.** *Chlidonias niger* (Linnaeus, 1758). (Photo: BDV).



**Figure 4.** Map of known wintering distribution of *Chlidonias niger* (Linnaeus, 1758) in South America (BirdLife International, Handbook of the Birds of the World 2018). Red circle is the first documented record for the Pantanal wetland and also for the Central region of Brazil. Black circles are secondary records of this species on the coast of Brazil and other places of South America (Dias et al. 2010; Quiroga et al. 2015; Gonsioroski; 2018; Franz et al. 2018; Wikiaves 2019).



**Figure 5.** **A.** *Aburria cumanensis* (Jacquin, 1784). **B.** *Egretta caerulea* (Linnaeus, 1758). **C.** *Calidris fuscicollis* (Vieillot, 1819). **D.** *Calidris melanotos* (Vieillot, 1819). **E.** *Phalaropus tricolor* (Vieillot, 1819). **F.** *Thectocercus acuticaudatus* (Vieillot, 1818). **G.** *Hirundo rustica* Linnaeus, 1758. **H.** *Petrochelidon pyrrhonota* (Vieillot, 1817). (Photos: BDV, except for *Aburria cumanensis*, photographed by AVBF).

individuals of *C. niger* were identified by their grey wings and back and white head and belly. The birds present with a dark crown and nape, with a black patch

behind the eye and an extensive dark patch on the side of the breast (Gochfeld et al. 2019).

This is the first record of *C. niger* for the Pantanal

wetland and the Central Region of Brazil (Fig. 4) The closest South American localities to our new record is Tres Pozos, Argentina (980 km) and Lagoa de Maricá, Rio de Janeiro, Brazil (1,675 km).

#### NEW RECORDS FOR ESTAÇÃO ECOLÓGICA DE TAIAMÃ

Order Galliformes  
Family Cracidae

##### *Aburria cumanensis* (Jacquin, 1784)

Figure 5A

**Material examined.** Tables 1, 2.

**Identification.** We generally observed this cracid in the forest of the EET. *Aburria cumanensis* differs from *Aburria kujubi* (Pelzeln, 1858) mainly by having a blue rather than red throat (Gwynne et al. 2010). We found both *Aburria* species in the EET, which had not been reported by Lopes et al. (2016).

Order Pelecaniformes  
Family Ardeidae

##### *Egretta caerulea* (Linnaeus, 1758)

Figure 5B

**Material examined.** Tables 1, 2.

**Identification.** We observed adult and immature individuals in mudslides and on beaches formed during the dry season. This heron can be readily identified by the predominantly blue colour of adults and white colour in the immature phase. The beak, tarsus, and toes are black (Sick 1997).

Order Accipitriformes  
Family Pandionidae

##### *Pandion haliaetus* (Linnaeus, 1758)

**Material examined.** Tables 1, 2.

**Identification.** This species was observed foraging on the Paraguay River during all hydrological periods, although it is classified as migratory. We identified this eagle by the combination of its white head and bottom and the presence of a black band behind the eye, which extends onto the upper part of the body (Sick 1997).

Family Accipitridae

##### *Leptodon cayanensis* (Latham, 1790)

**Material examined.** Tables 1, 2.

**Identification.** We observed this kite, usually alone, perched on the river's edge. We identified it by the small grey head, black back, and black tail with three whitish bars and a white tip (Gwynne et al. 2010).

##### *Ictinia mississippiensis* (Wilson, 1811)

**Material examined.** Tables 1, 2.

**Identification.** We recorded two individuals on different occasions in October 2016, which is during this species' migration to South America. The individuals were seen in flight near *Erythrina fusca* Monodominant Forest. This is a new record for Cáceres municipality. This

species differs from *I. plumbea* by its paler head, the lack of rufous primaries, the uniform dark tail, and the dark tarsus (Sick 1997; Gwynne et al. 2010).

Order Eurypygiformes  
Family Eurypygidae

##### *Eurypyga helias* (Pallas, 1781)

**Material examined.** Tables 1, 2.

**Identification.** This species was seen on the ground in polyspecific forests along the Paraguay River. We identified this species by the small head with a long, straight beak, and slender body with an unmistakable, complex colour (Gwynne et al. 2010).

Order Charadriiformes  
Family Scolopacidae

##### *Tringa flavipes* (Gmelin, 1789)

**Material examined.** Tables 1, 2.

**Identification.** We saw individuals foraging on mudflats and shoreline created during low water periods in Baía do Formoso, which forms part of the protection area buffer zone, and also in EET. This species differs from others of its genus by the slender, straight bill which is as long as the head, and the long yellow legs (Erize et al. 2006).

##### *Calidris fuscicollis* (Vieillot, 1819)

Figure 5C

**Material examined.** Tables 1, 2.

**Identification.** During the dry season, we observed *C. fuscicollis* on mudflats with other scolopacid species, both in Baía do Formoso, in the EET area, and buffer zone. This species presents with grey upper parts, a white eyebrow, white underparts, grey-striated chest and flanks, and a black bill and legs (Erize et al. 2006; Gwynne et al. 2010).

##### *Calidris melanotos* (Vieillot, 1819)

Figure 5D

**Material examined.** Tables 1, 2.

**Identification.** Individuals were observed foraging on mudflats with other scolopacid species. We identified *C. melanotos* by its brown, mixed with black-brown crown, neck, face, breast, and upperparts, white belly, and olive-yellow bill and legs (Erize et al. 2006; Van Gils et al. 2018).

##### *Phalaropus tricolor* (Vieillot, 1819)

Figure 5E

**Material examined.** Tables 1, 2.

**Identification.** Individuals were observed only in October 2016, when they were foraging with other scolopacids. This species was distinguished from all others on the mudflats by its white colour. This species was identified for by the grey upperparts, long slender bill, and white face, neck, and underparts (Erize et al. 2006).

Order Columbiformes  
Family Columbidae

##### *Columbina minuta* (Linnaeus, 1766)

**Material examined.** Tables 1, 2.

**Identification.** Small flocks of this species were observed around in the head office of the EET, where they were feeding on the ground. We identified this species by its dark-brown colour, with a bluish-grey head and pale reddish chest (Gwynne et al. 2010).

Order Caprimulgiformes

Family Caprimulgidae

***Nyctiprogne leucopyga* (Spix, 1825)**

**Material examined.** Tables 1, 2.

**Identification.** We observed this species during over-flight in the evening on the Paraguay River. We identified this species by the dark wing and black tail, which has a white bar midway (Erize et al. 2006).

Order Apodiformes

Family Apodidae

***Streptoprocne zonaris* (Shaw, 1796)**

**Material examined.** Tables 1, 2.

**Identification.** We recorded this species during over-flights on the Paraguay River. This is a new record for Cáceres municipality. This species presents a blackish plumage which is broken by a distinctive white collar on the neck (Chantler et al. 2019).

Order Piciformes

Family Ramphastidae

***Pteroglossus castanotis* Gould, 1834**

**Material examined.** Tables 1, 2.

**Identification.** We observed this species in various places in the EET, but it was especially common around the head office where it was eating fruits of *Sterculia apetala* (Jacq.) H. Karst. We identified this species by its song and coloration: the eyes are white and surrounded by blue skin; the back and wings are green; and the belly is yellow, crossed by a red band (Gwynne et al. 2010).

Order Falconiformes

Family Falconidae

***Falco ruficularis* Daudin, 1800**

**Material examined.** Tables 1, 2.

**Identification.** We saw this species only once, while it was preying on a nest of *Paroaria capitata* (d'Orbigny & Lafresnaye, 1837). This species was identified by the presence of a barred, white chest, with chestnut-rufous belly, thighs, and undertail-coverts (Gwynne et al. 2010).

Order Psittaciformes

Family Psittacidae

***Ara ararauna* (Linnaeus, 1758)**

**Material examined.** Tables 1, 2.

**Identification.** We observed these macaws only in flight on two occasions. We identified this species by its blue back, yellow belly, and white face (Gwynne et al. 2010).

***Thectocercus acuticaudatus* (Vieillot, 1818)**

Figure 5F

**Material examined.** Tables 1, 2.

**Identification.** This species was observed only during the dry season. Large flocks were feeding on the flowers and fruits of *E. fusca*. We identified this species by its blue forehead and predominantly green body (Gwynne et al. 2010).

***Pionus menstruus* (Linnaeus, 1766)**

**Material examined.** Tables 1, 2.

**Identification.** We recorded the voice of this species while in flight. The voice of this species differs from *Pionus maximiliani* (Kuhl, 1820) which also occurs in the EET.

Order Passeriformes

Family Dendrocolaptidae

***Xiphorhynchus guttatoides* (Lafresnaye, 1850)**

**Material examined.** Tables 1, 2.

**Identification.** This species was identified from a sound recording of its voice and by sight. It was in Polyspecific Forest of Shrubs and Pioneers.

Family Furnariidae

***Synallaxis hypospodia* Sclater, 1874**

**Material examined.** Tables 1, 2.

**Identification.** This species was identified from a sound recording of its voice. It was in Flooded Grassland and aquatic macrophytes. This is a new record for the Cáceres municipality.

Family Rynchocyclidae

***Tolmomyias sulphurens* (Spix, 1825)**

**Material examined.** Tables 1, 2.

**Identification.** This species was identified from a sound recording of its voice. It was in Polyspecific Forest of Shrubs and Pioneers.

Family Tyrannidae

***Elaenia spectabilis* Pelzeln, 1868**

**Material examined.** Tables 1, 2.

**Identification.** This species was identified from a sound recording of its voice. It was in Polyspecific Forest of Shrubs and Pioneers.

***Attila bolivianus* Lafresnaye, 1848**

**Material examined.** Tables 1, 2.

**Identification.** This species was identified from a sound recording of its voice. It was in Polyspecific Forest of Shrubs and Pioneers.

***Tyrannus savana* Daudin, 1802**

**Material examined.** Tables 1, 2.

**Identification.** This species was observed during migration. It was recognised by the characteristically long forked tail in flight (Gwynne et al. 2010).

Family Corvidae

***Cyanocorax cyanomelas* (Vieillot, 1818)**

**Material examined.** Tables 1, 2.

**Identification.** This species was identified from a sound recording of its voice. It was in Polyspecific Forest of Shrubs and Pioneers, next to the head office of the EET.

Family Hirundinidae

***Progne subis* (Linnaeus, 1758)**

**Material examined.** Tables 1, 2.

**Identification.** We observed a female of *P. subis* foraging on the banks of the river channel. The female of this species is dark at the top with some metallic blue luster and lighter underparts (Gwynne et al. 2010; Turner 2018).

***Riparia riparia* (Linnaeus, 1758)**

**Material examined.** Tables 1, 2.

**Identification.** This species was observed flying over the Paraguay River together with resident species like *Stelgidopteryx ruficollis* (Vieillot, 1817) and *Progne tapera* (Vieillot, 1817). We identified it as is a small swallow, with a uniform brown upper and somewhat darker wings. *Progne tapera* is similarly colored and patterned but is considerably larger (Ridgely and Tudor 2009).

***Hirundo rustica* Linnaeus, 1758**

Figure 5G

**Material examined.** Tables 1, 2.

**Identification.** This species was seen flying with individuals of *Petrochelidon pyrrhonota* (Vieillot, 1817); we observed many individuals foraging in the EET and buffer zone. We identified this species by its long, deeply forked tail and steely blue upper, with chestnut forehead and chestnut to buff throat (Ridgely and Tudor 2009).

***Petrochelidon pyrrhonota* (Vieillot, 1817)**

Figure 5H

**Material examined.** Tables 1, 2.

**Identification.** We observed this species in great flocks, foraging in the Flooded Grassland area. We observed a large number of foraging individuals in the EET and buffer zone. We identified this species by its square tail, dark blue upper, dirty white forehead and underneath, brown collar on the throat, face, and nape, and cinnamon upper tail (Ridgely and Tudor 2009).

Family Motacillidae

***Anthus lutescens* Pucheran, 1855**

**Material examined.** Tables 1, 2.

**Identification.** This species was identified from a sound recording of its voice made during the dry season in the flooded grassland.

**SPECIES IN IUCN RED LIST CATEGORY**

Order Galliformes

Family Cracidae

***Crax fasciolata* Spix, 1825**

**Material examined.** Tables 1, 2.

**Identification.** Species of Cracidae were observed in forests and around the EET head office. A couple was observed with two pups in November 2016 and again in 2017. In 2018, we saw a couple with three puppies. *Crax fasciolata* is considered globally Vulnerable (IUCN 2019). In Brazil, it is listed as Least Concern. This is the only species of *Crax* with a white tail tip. It has a grey bill and yellow cere. Males are mostly black, with white on the belly, while females have a bicolor crest, with a black back striped white and a cinnamon-coloured belly (Sick 1997; del Hoyo et al. 2018).

**OTHER NOTEWORTHY RECORDS**

We recorded other migratory and resident birds that are difficult to detect in the field due to their inconspicuous behaviour or occasional observations.

Order Charadriiformes

Family Scolopacidae

***Actitis macularius* (Linnaeus, 1766)**

Figure 2E

**Material examined.** Tables 1, 2.

**Identification.** One individual in eclipse plumage was seen foraging on a small mudflat on the shore of the Bracinho River. We identified this species by its brown upper, white belly and eyebrow, dark beak, and yellowish legs (Gwynne et al. 2010). We saw discreet spots on the belly, indicating its non-breeding status.

Order Pelecaniformes

Family Ardeidae

***Ixobrychus exilis* (Gmelin, 1789)**

Figure 2F

**Material examined.** Tables 1, 2.

**Identification.** This species was detected in all periods, mostly by voice. It was rarely seen but frequently heard. This species was observed making short flights and discreet landings among aquatic macrophytes (Pontederiaceae and Polygonaceae). *Ixobrychus exilis* has a characteristic voice and vocalises frequently. This species is a small, reddish-brown bittern, with two white stripes on the back. The male has a black cap and back, with yellowish stripes on the throat, while the female has a brown cap and back, with the darkest bands on the neck (Martínez-Vilata et al. 2018).

**GENERAL LIST OF EET BIRDS**

Field data, combined with data from the literature and museum specimens, yielded 278 bird species that occur in the EET. These species belong to 60 families in 24 orders (Table 2). Of these species, 160 were assigned to the primary list and 118 to the secondary list. None of the species recorded is endemic to the Pantanal biome. Of the species, 49 are migrant species, with 30 being partially migratory, 16 migratory, two vagrants, and one not



defined. In the general list, there are two globally Vulnerable species and one nationally Vulnerable species, *Penelope ochrogaster* Pelzeln, 1870.

For the tertiary list, seven species are included based on the known distribution their patterns (Table 3). We highlight the need for documented evidence of the species in the secondary list, especially for those whose documentation is unknown for the northwest region of the Pantanal, such as *Podilymbus podiceps* (Linnaeus, 1758), *Pardirallus nigricans* (Vieillot, 1819), and *Thamnophilus caerulescens* Vieillot, 1816, and the Nearctic migrants. *Tyrannus tyrannus* (Linnaeus, 1758) and *Sterna hirundo* Linnaeus, 1758.

## Discussion

The highly preserved and well-located Estação Ecológica de Taiamã, with its varied macrohabitats (Frota et al. 2017), harbours considerable bird richness, even though it is a predominantly open area. Nunes (2010) observed that this floodplain region shelters a greater number of species and individuals than the straighter and more human-disturbed segments of the Paraguay River. According to Lopes et al. (2016), Cáceres municipality, where EET is located, harbours the greatest number of bird species of all of Brazil's non-forest areas. This is due to the high diversity of habitats in the Cáceres municipality and because it lies at the contact zone between ecoregions, such as the Chiquitano Dry Forests and the Cerrado.

The Pantanal stands out in terms of species numbers when compared to other wetlands, such as the Florida Everglades or the Venezuelan Llanos. Our sampling comprised 35.56% of the Pantanal's avifauna (Nunes 2011), while the general list including the previously existing data comprised 47.76%. When systematic, non-systematic, LTER expeditions and secondary records are considered, we see the potential of the area to harbour high species richness, which may be due to its environmental heterogeneity regulated by hydrological variation.

Among all families, Tyrannidae tends to be abundant in the Pantanal region (Cintra and Yamashita 1990; Pinho et al. 2016; Vitorino et al. 2017) and represents one of the richest Neotropical bird families; the Tyraniidae is widely distributed in South and North America (Sick 1997). We highlight the representativeness of non-passerine families recorded in our study, and also reported in other Pantanal regions (Cintra and Yamashita 1990;

Antas et al. 2009; Nunes et al. 2010), due to the high species diversity in the families Ardeidae, Threskiornithidae, Anatidae, and Alcedinidae, among other non-passerines, which are associated with water bodies and open areas, such as found in the Pantanal.

Some key factors influencing variations in the frequency of bird occurrence in the EET include changes in the macrohabitats caused by the dry season and the presence of migratory species. Cintra and Yamashita (1990) also observed that seasonality and migratory movements affected the occurrence of bird species in northern Pantanal. In the southern Pantanal, Donatelli et al. (2017) found that even with a relatively stable abundance of bird species in different habitats over the years, there is a greater concentration of birds during the dry season, mainly due to lower water levels, which allow birds to explore more resources. In fact, the continuity of LTER in the EET can further promote information about fluctuations and enable the detection of other species.

The Pantanal is the final destination or stopover site for several migratory and nomad birds, which forage and breed there (Morrison et al. 2008; Oliveira et al. 2016). Ornithological inventories are extremely important, as they generate data on the occurrence and distribution patterns of bird species. In addition, monitoring can provide data on displacement and movement patterns throughout the year.

We highlight the record of *Chlidonias niger*, a migratory bird that breeds in North America and winters on the coast of Central America and South America. This species arrives back on the breeding grounds in April (Gochfeld et al. 2019), which is the same month as we observed an individual in the Pantanal. Our record adds information on the movements and habitat use of this species in Brazil. Until now, records in Brazil have been limited to coastal regions. Brazil is an important wintering destination for *C. niger*, which is usually observed between September and February (Somenzari et al. 2018). Individual birds in the Pantanal wetland are possibly vagrants, as seen in other intercontinental wetlands of South America, like in Chile and Argentina (Gochfeld et al. 2019). Nevertheless, the EET's wetlands ensure a very important stopover for this species.

Some resident species were only observed in certain periods, which strengthens the need for year-round monitoring to determine species' annual movements. Long-term studies are important not only for showing patterns of annual fluctuations but also to detect the occurrence

**Table 3.** Tertiary list of birds recorded at Estação Ecológica de Taiamã, Mato Grosso Pantanal, Brazil. Birds without a documented evidence, whose occurrence is questionable, invalid, or non-existent in the Pantanal wetlands.

*Crypturellus noctivagus* (Wied, 1820): Listed by Brazil (2017). No context given. Distribution unknown to the Pantanal wetlands.

*Aramides saracura* (Spix, 1825): Listed by Brazil (2017). No context given. Distribution unknown to the Pantanal wetlands.

*Gallinula angulata* Sundevall, 1850: Listed by Brazil (2017). No context given. Distribution unknown to the Pantanal wetlands.

*Celeus flavescens* (Gmelin, 1788): Listed by Brazil (2017). No context given. Distribution unknown to the Pantanal wetlands.

*Brotogeris versicolurus* (Statius Müller, 1776): Listed by Brazil (2017). No context given. Distribution unknown to the Pantanal wetlands. Likely use of old nomenclature for *Brotogeris chiriri*.

*Thamnophilus palliatus* (Lichtenstein, 1823): Listed by Brazil (2017). No context given. Distribution unknown to the Pantanal wetlands.

*Conopophaga lineata* (Wied, 1831): Listed by Brazil (2017). No context given. Distribution unknown to the Pantanal wetlands.

of species, which may be associated with the availability of food resources (Alves 2007). As for migrant species of the Hirundinidae family, we associate their occurrence with an abundance of their food resources, the insect flock, in the Pantanal wetland.

During the dry season, water levels are low, and especially in the buffer zone, this provides limniculous habitats for migrant scolopacids to forage, whereas the formation of beaches offers habitat for the nesting of colonial species such as *Rynchops niger*, as also observed by Antas et al. (2016) on the Cuiabá River.

The habitat occupation patterns differ among migratory species (Signor and Pinho 2011) and the abundance of residents is related to the use of a greater variety of Pantanal habitats (Figueira et al. 2011). The various macrohabitats such Aquatic-Terrestrial Transition Zones, Marshes, and Permanently Flooded Areas in the of the EET and buffer zone are due to the hydric dynamics of the Paraguay River and favour a greater availability of habitats and resources for bird species.

Migratory birds depend on habitats and food resources in breeding and wintering grounds that are often thousands of kilometres apart (Oliveira et al. 2016). If these habitats and resources are not protected, the birds are at risk of great population losses. According to Bird Life International (2018), around 40% of migratory bird species in decline.

Da Silva et al. (2015) and Tomas et al. (2019) stated that the diverse impacts that take place in Pantanal wetland directly affect ecosystem dynamics caused by the hydrological regime and landscape changes, such as hydroelectric power plants and the Paraguay River Waterway (Hydrovia). Cintra (2011) emphasised that the protection and conservation of rivers, baías (floodplain ponds), and lake beaches, as habitats adjacent, are fundamental for maintaining bird diversity. Thus, conservation actions must protect of the Pantanal wetlands and also of the surrounding plateau (Pinho et al. 2017).

The EET harbours bird species typical of wetlands as well as some savanna species. There are no species endemic to the Pantanal biome, but species directly associated with or dependent on these wetlands are well represented, such as *Cercomacra melanaria* (Ménétrières, 1835) and *Synallaxis albilora* Pelzeln, 1856. The importance of bird surveys in the Pantanal biome is, thus, underscored, especially given the difficulty in detecting resident and migratory species in the EET.

Environmental heterogeneity of the EET may be observed by species richness and composition. For example, some birds inhabiting flooded grasslands, such as species of the genus *Sporophila*; from floating macrophyte mats, like *Amblyramphus holosericeus*; from aquatic areas, like *Heliornis fulica*; from mudflats, such as species in the family Scolopacidae; from open areas, like *Ardea cocoi* Linnaeus, 1766 and *Ciconia maguari* (Gmelin, 1789); and from forests, such as *Campylorhamphus trochilrostris* (Lichtenstein, 1820), *Trogon curucui*

Linnaeus, 1766, *Procacicus solitarius* (Vieillot, 1816), and *Cercomacra melanaria*.

The occurrence of some as migrant and vagrant species in the EET shows that this area is very significant for maintaining the ecosystem and its biodiversity. Documentation of the bird species is necessary to support scientific knowledge of this wetland. Expanding the boundaries of this protected area further may promote its conservation, as the EET is important as a stop-over, feeding and breeding ground, and wintering site for migratory birds. Long-term monitoring in the EET is needed for an understanding of the ecology of these bird species in this protected area and for improved management and conservation of the Pantanal wetland.

## Acknowledgements

We wish to thank CAPES for a scholarship granted to AVBF and BDV; CNPq for support in Long-Term Ecological Research in the Pantanal wetland “Ecological Dynamics of the Upper Paraguay River floodplain” (no. 441563/2016-3); the Programa de Pós-graduação em Ciências Ambientais of Universidade do Estado de Mato Grosso, Cáceres; SISBIO/IBAMA for authorization to scientific activities in this protected area; Instituto Chico Mendes de Conservação da Biodiversidade and Daniel Kantek, Thadeu Deluque, Selma Myiazaki and Rogério Costa of the Estação Ecológica de Taiamã staff for their logistic support; Alex Lees and Vitor Piacentini for helping to determine of *Chlidonias niger* species.

## Authors' Contributions

AVBF project design, data collection, identification of species, compilation, and review of occurrence records and preparation of the manuscript. BDV project design, data collection, species identification, and review of the manuscript. CJS project design and review of the manuscript. SKI and JRN review of the manuscript.

## References

- Alves MAS (2007) Sistemas de migrações de aves em ambientes terrestres no Brasil: exemplos, lacunas e propostas para o avanço do conhecimento. *Revista Brasileira de Ornitologia* 15 (2): 231–238.
- Antas PTZ, Carrara LA, Ubaid FK, Oliveira-Junior SB, Ferreira LP (2016) Aves coloniais da Reserva Particular do Patrimônio Natural Sesc Pantanal. *Conhecendo o Pantanal* 10. SESC Departamento Nacional. Rio de Janeiro, 236 pp.
- Antas PTZ, Palo-Junior H (2009) *Pantanal: guia de aves*. 2ª edição. SESC Departamento Nacional, Rio de Janeiro, 249 pp.
- BirdLife International and Handbook of the Birds of the World (2018) Bird species distribution maps of the world. Version 2018.1. <http://datazone.birdlife.org/species/requestdis>. Accessed on: 2019-05-31.
- BirdLife International (2019) <http://www.birdlife.org/worldwide/programmes/migratory-birds-and-flyways>. Accessed on: 2019-05-20.
- Blondel J, Ferry C, Frochot B (1970) La method des indices ponctuels d'abondance (IPA) ou des relevés d'avifaune par stations d'écoute. *Alauda* 38 (1): 55–71.
- Brasil (2000) Lei Federal nº 9985 de 18 de julho de 2000 que institui o

- Sistema Nacional de Unidades de Conservação da Natureza. Ministério do Meio Ambiente, Brasília. [http://www.planalto.gov.br/ccivil\\_03/leis/19985.htm](http://www.planalto.gov.br/ccivil_03/leis/19985.htm). Accessed on 2019-03-29.
- Brasil (2017) Plano de Manejo da Estação Ecológica de Taiamã. Instituto Chico Mendes de Conservação da Biodiversidade, Ministério do Meio Ambiente, Brasília, 174 pp. [http://www.icmbio.gov.br/portal/images/stories/plano-de-manejo/plano\\_de\\_manejo\\_esec\\_taiama\\_vs\\_070617.pdf](http://www.icmbio.gov.br/portal/images/stories/plano-de-manejo/plano_de_manejo_esec_taiama_vs_070617.pdf). Accessed on: 2019-03-29.
- Brasil (2018) Livro vermelho da Fauna Brasileira ameaçada de extinção. Instituto Chico Mendes de Conservação da Biodiversidade, Ministério do Meio Ambiente, Brasília, 712 pp. [http://www.icmbio.gov.br/portal/images/stories/comunicacao/publicacoes/publicacoes-diversas/livro\\_vermelho\\_2018\\_vol3.pdf](http://www.icmbio.gov.br/portal/images/stories/comunicacao/publicacoes/publicacoes-diversas/livro_vermelho_2018_vol3.pdf). Accessed on: 2019-03-29.
- Brasil (2019) Ministério do Meio Ambiente. Sítios Ramsar. <https://www.mma.gov.br/biodiversidade/biodiversidade-aquatica/zonas-umidas-convencao-de-ramsar/sitios-ramsar.html>. Accessed on: 2019-03-29.
- Carlos CJ, Straube FC, Pacheco JF (2010) Conceitos e definições sobre documentação de registros ornitológicos e critérios para a elaboração de listas de aves para os estados brasileiros. *Revista Brasileira de Ornitologia* 18 (4): 355–361.
- Cintra R (2011) Pantanal de Poconé. In: Valente RM, Silva JMC, Straube FC, Nascimento JLX (Eds) Conservação de aves migratórias neárticas no Brasil. Conservação Internacional, Belém, 193–198.
- Cintra R, Yamashita C (1990) Habitats, abundância e ocorrência das espécies de aves do Pantanal de Poconé, Mato Grosso, Brasil. *Papéis Avulsos de Zoologia* 37 (1): 1–21.
- Chantler P, Kirwan, GM, Juana E (2019) White-collared Swift (*Streptoprocne zonaris*). In: Del Hoyo J, Elliott A, Sargatal J, Christie DA, De Juana E (Eds) Handbook of the birds of the world alive. Lynx Edicions, Barcelona. <https://www.hbw.com/node/55260>. Accessed on: 2019-03-29.
- Da Silva CJ, Sousa KNS, Ikeda-Castrillon SK, Lopes CRAS, Nunes JRS, Carniello MC, Mariotti PR, Lazaro WL, Morini A, Zago BW, Façanha CL, Albernaz-Silveira R, Loureiro E, Viana IG, Oliveira RF, Cruz WJA, Arruda JC, Sander NL, Freitas Junior DS, Pinto VR, Lima AC, Jongman RHG (2015) Biodiversity and its drivers and pressures of change in the wetlands of the Upper Paraguay–Guaporé Ecotone, Mato Grosso (Brazil). *Land Use Policy* 47: 163–178. <https://doi.org/10.1016/j.landusepol.2015.04.004>
- Dias RA, Agne CE, Gianuca D, Gianuca A, Barcellos-Silveira A, Bugoni L (2010) New records, distribution and status of six seabird species in Brazil. *Iheringia, Série Zoologia* 100 (4): 379–390. <https://doi.org/10.1590/S0073-47212010000400013>
- Del Hoyo J, Elliott A, Sargatal J, Christie DA, De Juana E (2017) Handbook of the birds of the world alive. Lynx Editions, Barcelona. <https://www.hbw.com>. Accessed on: 2018-01-08.
- Del Hoyo J, Kirwan GM, Sharpe CJ (2018) Bare-faced Curassow (*Crax fasciolata*). In: del Hoyo J, Elliott A, Sargatal, J, Christie DA, de Juana E (Eds) Handbook of the birds of the world alive. Lynx Edicions, Barcelona. <https://www.hbw.com/node/53316>. Accessed on: 2018-01-08.
- Donatelli RJ, Eaton DP, Cardoso GS, Vianna RM, Gerotti RW, Rodrigues FG, Martins RM (2017) Temporal and spatial variation of richness and abundance of the community of birds in the Pantanal wetlands of Nhecolândia (Mato Grosso do Sul, Brazil). *Revista Biologia Tropical* 65 (4): 1358–1380. <https://doi.org/10.15517/rbt.v65i4.27729>
- Franz I, Agne CE, Bencke GA, Bugoni L, Dias RA (2018) Four decades after Belton: a review of records and evidences on the avifauna of Rio Grande do Sul, Brazil. *Iheringia Série Zoologia* 108: 1–38. <https://doi.org/10.1590/1678-4766e2018005>
- Frota AVB, Ikeda-Castrillon SK, Kanteck DLZ, Silva CJ (2017). Macrohabitats da Estação Ecológica de Taiamã, no contexto da Área Úmida Pantanal mato-grossense, Brasil. *Boletim do Museu Paraense Emílio Goeldi, Ciências Naturais* 12 (2): 263–280.
- Figueira JEC, Mourão FA, Coelho AS (2011) Habitat heterogeneity and climatic seasonality structure the avifauna trophic guilds in the Brazilian Pantanal wetland. *Canadian Journal of Zoology* 89: 1206–1213. <https://doi.org/10.1139/z11-099>
- Gochfeld M, Burger J, Garcia EFJ (2019) Black Tern (*Chlidonias niger*). In: Del Hoyo J, Elliott A, Sargatal J, Christie DA, De Juana E (Eds) Handbook of the birds of the world alive. Lynx Edicions, Barcelona. <https://www.hbw.com/node/54047>. Accessed on: 2019-05-30.
- Gonsioroski G (2014) Primeiro registro documentado de *Leucophaeus pipixcan* e novos registros de *Stercorarius parasiticus* e *Chlidonias niger* (Charadriiformes) no estado do Maranhão. *Atualidades Ornitológicas* 180: 14–15.
- Gwynne JA, Ridgely RS, Tudor G, Argel M (2010) Aves do Brasil: Pantanal & Cerrado. Horizonte, São Paulo, 322 pp.
- Erize F, Mata JRR, Rumboll M (2006) Birds of South America, non-passerines: heas to woodpeckers. Princeton University Press, Princeton, 385 pp.
- IUCN (2019) The IUCN Red List of threatened species. International Union for Conservation of Nature, Gland. <http://www.iucnredlist.org>. Accessed on: 2019-05-20.
- Junk WJ, Bayley PB, Sparks RE (1989) The flood pulse concept in river–floodplain systems. *Canadian Special Publication of Fisheries and Aquatic Sciences* 106: 110–127.
- Junk WJ, Cunha CN, Wantzen KM, Petermann P, Strüssmann C, Marques MI, Adis J (2006) Biodiversity and its conservation in the Pantanal of Mato Grosso, Brazil. *Aquatic Sciences* 68: 278–309. <https://doi.org/10.1007/s00027-006-0851-4>
- Junk WJ, Da Silva CJ (1996) O conceito do pulso de inundação e suas implicações para o Pantanal de Mato Grosso. In: Dantas M, Catto JB, Resende EK (Eds) II Simpósio sobre Recursos Naturais e Socioeconômicos do Pantanal: manejo e conservação. EMBRAPA, Corumbá, 17–28.
- Kanteck DLZ, Onuma SSM (2013) Primeiro registro documentado da gaivota-de-franklin *Leucophaeus pipixcan* Wagler, 1831 para o bioma Pantanal, Brasil. *Ornithologia* 6 (1): 106–108.
- Kotteck M, Grieser J, Beck C, Rudolf B, Rubel F (2006) World map of the Köppen–Geiger climate classification updated. *Meteorologische Zeitschrift* 15 (3): 259–263. <https://doi.org/10.1127/0941-2948/2006/0130>
- Lopes LE, Pinho JB, Ortiz A, Evangelista MM, Silveira LF, Schunck F, Devely PF (2016) Birds from Cáceres, Mato Grosso: the highest species richness ever recorded in a Brazilian non-forest region. *Revista Brasileira de Ornitologia* 24 (2): 137–167.
- Martínez-Vilalta A, Motis A, Kirwan GM (2018) Least Bittern (*Ixobrychus exilis*). In: Del Hoyo J, Elliott A, Sargatal J, Christie DA, De Juana E (Eds) Handbook of the birds of the world alive. Lynx Edicions, Barcelona. <https://www.hbw.com/node/52721>. Accessed on: 2018-01-08.
- Morrison RIG, Serrano IL, Antas PTZ, Ross K (2008) Aves migratórias no Pantanal: distribuição de aves limícolas neárticas e outras espécies aquáticas no Pantanal. WWF-Brasil, Brasília, 100 pp.
- Nunes AP (2011) Quantas espécies de aves ocorrem no Pantanal brasileiro? *Atualidades Ornitológicas* 160: 45–54.
- Nunes AP, Tizianel FAT, Melo AV, Nascimento V, Machado N (2010) Aves da Estrada Parque Pantanal, Corumbá, Mato Grosso do Sul, Brasil. *Atualidades Ornitológicas On-line* 156: 33–47.
- Nunes JRS (2010) Avifauna do Rio Paraguai, Pantanal de Cáceres, Mato Grosso. PhD dissertation, Universidade Federal de São Carlos, São Carlos, 259 pp.
- Nunes AP, Straube FC, Laps RR, Posso SR (2017) Checklist das aves do Mato Grosso do Sul, Brasil. *Iheringia Série Zoologia* 107: 1–19. <https://doi.org/10.1590/1678-4766e2017154>
- Oliveira AC, Barbosa AEA, Sousa AEA, Lugarini C, Lima DM, Nascimento JLX, Souza MA, Somenzari M, Souza NA, Serafini PP, Amaral PP, Rossato RM, Medeiros RCS (2016) Relatório anual de rotas e áreas de concentração de aves migratórias no Brasil. CEMAVE/ICMBio, Cabedelo, 63 pp.

- Piacentini VQ, Aleixo A, Agne CE, Maurício GN, Pacheco JF, Bravo GA, Brito GRR, Naka LN, Olmos F, Posso S, Silveira LF, Betini GS, Carrano E, Franz I, Lees AC, Lima LM, Pioli D, Schunck F, Amaral FR, Bencke GA, Cohn-Haft M, Figueiredo LFA, Straube FC, Cesari E (2015) Annotated checklist of the birds of Brazil by the Brazilian Ornithological Records Committee. *Revista Brasileira de Ornitologia* 23 (2): 91–298.
- Pinho JB, Lopes LE, Marini MA (2016) Birds from the Pirizal region, Pantanal of Poconé, Mato Grosso, Brazil. *Revista Brasileira de Ornitologia* 24 (3): 267–285.
- Pinho JB, Aragona M, Hakamada KYP, Marini MA (2017) Migrations patterns and seasonal forest use by birds in the Brazilian Pantanal. *Bird Conservation International* 27 (3): 371–387. <https://doi.org/10.1017/S0959270916000290>
- Quiroga OB, CoriaOR, Gómez N, Chimino G, Jorge F, Rojas LM, Llugdar JE, Alcalde M (2015) El Gaviotín Negro (*Chlidonias niger*) yel Burrito Común (*Laterallus melanophaius*) en Santiago Del Estero, Argentina. *Nuestras Aves* 60: 59–60.
- Ridgely RS, Tudor G (2009) Field guide to the songbirds of South America: the passerines. University of Texas Press, Austin, 751 pp.
- Sick H (1997) Ornitologia brasileira. Nova Fronteira, Rio de Janeiro, 912 pp.
- Signor CA, Pinho JB (2011) Spatial diversity patterns of birds in a vegetation mosaic of the Pantanal, Mato Grosso, Brazil. *Zoologia* 28 (6): 725–738. <https://doi.org/10.1590/S1984-46702011000600005>
- Silveira LF, Beisiegel BDM, Curcio FF, Valdujo PH, Dixo M, Verdade VK, Mattox GMT, Cunningham PTM (2010) What use do fauna inventories serve? *Estudos Avançados* 24 (68): 173–207.
- Somenzari M, do Amaral PP, Cueto VR, de Camargo Guaraldo A, Jahn AE, Lima DM, Lugarini C, Machado CG, Martínez J, do Nascimento JLX, Pacheco JF, Paludo D, Prestes NP, Serafini PP, Silveira LF, de Sousa EBA, Alves de Sousa N, de Souza MA, Telino-Junior WR, Whitney BM (2018) An overview of migratory birds in Brazil. *Papéis Avulsos de Zoologia* 58(3): 1–66. <https://doi.org/10.11606/1807-0205/2018.58.03>
- Tomas WM, Roque FO, Morato RG, Medici PE, Chiaravalloti RM, Tortato FR, Penha JMF, Nzo TJ, Garcia, LC, Lourival RFF, Girard P, Albuquerque IR, Almeida-Gomes M, Andrade MHS, Araujo AS, Araujo C, Arruda EC, Battiolola LD, Benites M, Assunção VA, Bolzan FP, Boock JC, Bortolotto IM, Brasil MS, Camilo AR, Campos Z, Carniello MA, Catella AC, Cheida CC, Crawshaw Jr PG, Crispim SMA, Damasceno Junior GA, Desbiez ALJ, Dias FA, Eaton DP, Faggioni GP, Farinaccio MA, Fernandes JFA, Ferreira VL, Fischer EA, Frago CE, Freitas GO, Galvani F, Garcia AS, Garcia CM, Gracioli G, Guariento RD, Guedes NMR, Guerra A, Herrera HM, Hoogesteijn R, Ikeda-Castrillon SK, Juliano RS, Kantek DLZ, Keuroghlian A, Lacerda ACR, Lacerda ALR, Landeiro VL, Laps RR, Layme V, Leimgruber P, Rocha FL, Mamede S, Marques DKS, Marques MI, Mourão GM, Moraes RN, Moreira TA, Nicola RD, Nogueira DG, Nunes AP, Nunes da Cunha C, Oliveira MD, Oliveira MR, Paggi GM, Pellegrin AO, Pereira GMF, Peres IAHSF, Pinho JB, Pinto JOP, Pott A, Provete DB, Reis VDA, Reis RK, Renaud P, Ribeiro DB, Rosseto OC, Sabino J, Rumiz D, Salis SM, Santana DJ, Santos AS, Sartori AL, Sato M, Schuchmann KL, Scremin-Dias E, Sigrist MR, Silva A, Da Silva CJ, Siqueira AL, Soriano BMA, Sousa LM, Souza FL, Strussmann C, Sugai LSM, Tocantins N, Urbanetz C, Valente-Neto F, Viana DP, Yanosky A, Junk WJ (2019) Sustainability agenda for the Pantanal Wetland: perspectives on a collaborative interface for science, policy, and decision-making. *Tropical Conservation Science* 12: 1–30. <https://doi.org/10.1177/1940082919872634>
- Tubelis DP, Tomas WM (2003) Bird species of the Pantanal wetland, Brazil. *Ararajuba* 11 (1): 5–37.
- Turner A (2018) Purple Martin (*Progne subis*). In: Del Hoyo J, Elliott A, Sargatal J, Christie DA, De Juana E (Eds) *Handbook of the birds of the world alive*. Lynx Edicions, Barcelona. <https://www.hbw.com/node/57709>. Accessed on: 2018-01-15.
- Van Gils J, Wiersma P, Christie DA, Kirwan GM (2018) Pectoral Sandpiper (*Calidris melanotos*). In: Del Hoyo J, Elliott A, Sargatal J, Christie DA, De Juana E (Eds) *Handbook of the birds of the world alive*. Lynx Edicions, Barcelona. <https://www.hbw.com/node/53933>. Accessed on: 2018-01-15.
- Veloso HP, Rangel-Filho ALT, Lima JCA (1991) Classificação da vegetação brasileira adaptada a um sistema universal. IBGE, Rio de Janeiro, 124 pp.
- Vitorino BD, Frota AVB, Angelo M, Nunes JRS (2017) Avifauna associada a duas áreas de nascentes no Assentamento Laranjeira I, Província Serrana, Cáceres-MT. In: Ikeda-Castrillon S, Puhl JI, Morais FF, Morini-Lopes AAET (Eds) *Escassez hídrica e restauração ecológica no Pantanal: recuperação das nascentes e fragmentos de mata ciliar do córrego no Assentamento Laranjeira I e mobilização para conservação dos recursos hídricos no Pantanal mato-grossense*. 1 ed. Carlini & Caniato Editorial, Cuiabá, 153–167.
- Vitorino BD, Frota AVB, Ikeda-Castrillon SI, Nunes JRS (2018) Birds of Estação Ecológica da Serra das Araras, state of Mato Grosso, Brazil: additions and review. *Check List* 14 (5): 893–922. <https://doi.org/10.15560/14.5.893>
- Wantzen KM, Drago E, Da Silva CJ (2005) Aquatic habitats of the Upper Paraguay river-floodplain-system and parts of the Pantanal (Brazil). *Ecohydrology and Hydrobiology* 5(2): 107–126.
- Wikiaves (2019) <http://www.wikiaves.com/>. Accessed on: 2019-05-31.

## Appendix

**Table S1.** Survey sites, site codes, and points, with their geographic positions and macrohabitats.

Hydrological cycle	Site name	Site code	Point	Latitude	Longitude	Macrohabitats
2015–2016	Station 1	S1	MF1P1	16°49.36'S	057°32.48'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			MF1P2	16°49.24'S	057°32.45'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			MF1P3	16°49.17'S	057°32.36'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			MF1P4	16°49.28'S	057°32.30'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			MF1P5	16°49.33'S	057°32.19'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			MF1P6	16°49.27'S	057°32.06'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			MF1P7	16°49.20'S	057°31.97'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			MF1P8	16°49.19'S	057°31.84'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
Station 2	S2	MF2P1	16°52.27'S	057°32.06'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels	
		MF2P2	16°52.15'S	057°32.13'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels	
		MF2P3	16°52.04'S	057°32.19'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels	
		MF2P4	16°51.95'S	057°32.26'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels	
		MF2P5	16°51.86'S	057°32.35'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels	
		MF2P6	16°51.82'S	057°32.46'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels	

Hydrological cycle	Site name	Site code	Point	Latitude	Longitude	Macrohabitats
2015–2016	Station 2	S2	MF2P7	16°51.67'S	057°33.42'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels
			MF2P8	16°51.62'S	057°33.54'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / floating islands and meadows / river channels
	Station 3	S3	PF1P1	16°51.89'S	057°33.17'W	Shrubland and pioneer forest / river channels
			PF1P2	16°51.95'S	057°33.06'W	Shrubland and pioneer forest / river channels
			PF1P3	16°51.94'S	057°32.94'W	Shrubland and pioneer forest / river channels
			PF1P4	16°51.92'S	057°32.88'W	Shrubland and pioneer forest / river channels
			PF1P5	16°51.89'S	057°32.69'W	Shrubland and pioneer forest / river channels
			PF1P6	16°50.86'S	057°35.17'W	Shrubland and pioneer forest / river channels
			PF1P7	16°50.95'S	057°35.11'W	Shrubland and pioneer forest / river channels
			PF1P8	16°51.04'S	057°35.02'W	Shrubland and pioneer forest / river channels
	Station 4	S4	PF2P1	16°52.18'S	057°26.47'W	Shrubland and pioneer forest / river channels
			PF2P2	16°52.29'S	057°26.42'W	Shrubland and pioneer forest / river channels
			PF2P3	16°52.38'S	057°26.34'W	Shrubland and pioneer forest / river channels
			PF2P4	16°52.48'S	057°26.29'W	Shrubland and pioneer forest / river channels
			PF2P5	16°52.56'S	057°26.19'W	Shrubland and pioneer forest / river channels
			PF2P6	16°52.56'S	057°26.07'W	Shrubland and pioneer forest / river channels
			PF2P7	16°52.58'S	057°25.96'W	Shrubland and pioneer forest / river channels
			PF2P8	16°52.62'S	057°25.84'W	Shrubland and pioneer forest / river channels
	Station 5	S5	FG1P1	16°49.71'S	057°38.11'W	Flooded grassland / floating islands and meadows / river and floodplain channels
			FG1P2	16°49.82'S	057°38.09'W	Flooded grassland / floating islands and meadows / river and floodplain channels
			FG1P3	16°49.94'S	057°38.07'W	Flooded grassland / floating islands and meadows / river and floodplain channels
			FG1P4	16°50.04'S	057°38.14'W	Flooded grassland / floating islands and meadows / river and floodplain channels
			FG1P5	16°50.15'S	057°38.18'W	Flooded grassland / floating islands and meadows / river and floodplain channels
			FG1P6	16°50.23'S	057°38.27'W	Flooded grassland / floating islands and meadows / river and floodplain channels
			FG1P7	16°50.30'S	057°38.35'W	Flooded grassland / floating islands and meadows / river and floodplain channels
			FG1P8	16°50.25'S	057°38.45'W	Flooded grassland / floating islands and meadows / river and floodplain channels
	Station 6	S6	FG2P1	16°52.75'S	057°24.61'W	Flooded grassland / floating islands and meadows / lakes / river channels
			FG2P2	16°52.72'S	057°24.51'W	Flooded grassland / floating islands and meadows / lakes / river channels
			FG2P3	16°52.63'S	057°24.47'W	Flooded grassland / floating islands and meadows / lakes / river channels
			FG2P4	16°52.51'S	057°24.45'W	Flooded grassland / floating islands and meadows / lakes / river channels
			FG2P5	16°52.45'S	057°24.31'W	Flooded grassland / floating islands and meadows / lakes / river channels
			FG2P6	16°52.48'S	057°24.18'W	Flooded grassland / floating islands and meadows / lakes / river channels
			FG2P7	16°52.38'S	057°24.14'W	Flooded grassland / floating islands and meadows / lakes / river channels
			FG2P8	16°52.26'S	057°24.19'W	Flooded grassland / floating islands and meadows / lakes / river channels
2017–2018	Track 1	T1	T1P1	16°49.32'S	057°32.24'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P2	16°49.18'S	057°31.91'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P3	16°49.27'S	057°31.68'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P4	16°49.49'S	057°31.40'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P5	16°49.53'S	057°31.13'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P6	16°49.44'S	057°30.80'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P7	16°49.33'S	057°30.61'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P8	16°49.48'S	057°30.38'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P9	16°49.68'S	057°30.18'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
			T1P10	16°49.55'S	057°29.93'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / aquatic macrophytes / river channels
	Track 2	T2	T2P1	16°51.77'S	057°26.14'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P2	16°52.07'S	057°26.33'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P3	16°52.46'S	057°26.28'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P4	16°52.57'S	057°26.00'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P5	16°52.64'S	057°25.69'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P6	16°52.57'S	057°25.35'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P7	16°52.45'S	057°25.14'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P8	16°52.70'S	057°25.07'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P9	16°52.98'S	057°25.06'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
			T2P10	16°52.94'S	057°24.68'W	Shrubland and pioneer forest / aquatic macrophytes / river channels
	Track 3	T3	T3P1	16°53.39'S	057°29.62'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels
			T3P2	16°53.67'S	057°29.45'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels
			T3P3	16°54.03'S	057°29.36'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels
			T3P4	16°53.98'S	057°28.93'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels

Hydrological cycle	Site name	Site code	Point	Latitude	Longitude	Macrohabitats		
2017–2018	Track 3	T3	T3P5	16°54.06'S	057°28.63'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels		
			T3P6	16°54.66'S	057°28.92'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels		
			T3P7	16°54.59'S	057°28.33'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels		
			T3P8	16°54.92'S	057°28.26'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels		
			T3P9	16°55.03'S	057°28.00'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels		
			T3P10	16°55.01'S	057°27.72'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels		
			Track 4	T4	T4P1	16°51.69'S	057°33.50'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels
					T4P2	16°51.81'S	057°33.30'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels
					T4P3	16°51.93'S	057°33.00'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels
					T4P4	16°51.91'S	057°32.68'W	Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels
T4P5	16°51.86'S	057°32.41'W			Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels			
T4P6	16°52.07'S	057°32.23'W			Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels			
T4P7	16°52.33'S	057°32.04'W			Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels			
T4P8	16°51.99'S	057°31.96'W			Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels			
T4P9	16°51.80'S	057°31.86'W			Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels			
T4P10	16°51.64'S	057°31.64'W			Monospecific forest with <i>Erythrina fusca</i> Lour. / shrubland and pioneer forest / flooded grasslands / aquatic macrophytes / floating islands and meadows / river channels			
Track 5	T5	T5P1	16°49.00'S	057°39.43'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P2	16°49.22'S	057°39.59'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P3	16°49.54'S	057°39.55'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P4	16°49.81'S	057°39.64'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P5	16°50.06'S	057°39.70'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P6	16°50.33'S	057°39.83'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P7	16°50.60'S	057°39.63'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P8	16°50.87'S	057°39.56'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P9	16°51.04'S	057°39.28'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			
		T5P10	16°51.19'S	057°39.51'W	Flooded grassland / aquatic macrophytes / floating islands and meadows / lakes / river and floodplain channels			